

w o r k i n g
p a p e r

14 13

**Evaluating Conditional Forecasts from
Vector Autoregressions**

Todd E. Clark and Michael W. McCracken



FEDERAL RESERVE BANK OF CLEVELAND

Working papers of the Federal Reserve Bank of Cleveland are preliminary materials circulated to stimulate discussion and critical comment on research in progress. They may not have been subject to the formal editorial review accorded official Federal Reserve Bank of Cleveland publications. The views stated herein are those of the authors and are not necessarily those of the Federal Reserve Bank of Cleveland or of the Board of Governors of the Federal Reserve System.

Working papers are available at:

www.clevelandfed.org/research.

Evaluating Conditional Forecasts from Vector Autoregressions

Todd E. Clark and Michael W. McCracken

Many forecasts are conditional in nature. For example, a number of central banks routinely report forecasts conditional on particular paths of policy instruments. Even though conditional forecasting is common, there has been little work on methods for evaluating conditional forecasts. This paper provides analytical, Monte Carlo, and empirical evidence on tests of predictive ability for conditional forecasts from estimated models. In the empirical analysis, we consider forecasts of growth, unemployment, and inflation from a VAR, based on conditions on the short-term interest rate. Throughout the analysis, we focus on tests of bias, efficiency, and equal accuracy applied to conditional forecasts from VAR models.

JEL Nos.: C53, C52, C12, C32.

Keywords: Prediction, forecasting, out-of-sample.

Suggested citation: Todd E. Clark and Michael W. McCracken, 2014. "Evaluating Conditional Forecasts from Vector Autoregressions," Federal Reserve Bank of Cleveland, working paper no. 14-13.

Todd E. Clark is at the Federal Reserve Bank of Cleveland (P.O. Box 6387; Cleveland, OH 44101; todd.clark@clev.frb.org). Michael W. McCracken is at the Federal Reserve Bank of St. Louis (P.O. Box 442; St. Louis, MO 63166; michael.w.mccracken@stls.frb.org). The authors gratefully acknowledge helpful comments from Graham Elliott, Ana Galvao, Silvia Goncalves, James Nason, Frank Schorfheide, Dan Waggoner, Mark Watson, and participants at the 2014 NBER Summer Institute, 2014 Barcelona GSE Summer Forum on Time Series Analysis, the IAAE 2014 Annual Conference, and the 2014 EABCN conference on Judgment and Combination in Forecasting and Policy Models. The views expressed herein are solely those of the authors and do not necessarily reflect the views of the Federal Reserve Bank of Cleveland, Federal Reserve Bank of St. Louis, Federal Reserve System, or any of its staff.

1 Introduction

Since the seminal work of Litterman (1986), vector autoregressions (VARs) have been widely known to be useful for out-of-sample forecasting. In many applications, the forecasts of interest are unconditional. Clark and McCracken (2013) review methods for evaluating such forecasts.

VARs are also commonly used to construct conditional forecasts. In such applications, the models are used to predict variables such as GDP growth and inflation conditional on, e.g., an assumed path of monetary policy variables or an assumed path of oil prices. Examples of VAR forecasts conditional on policy paths include Sims (1982), Doan, Litterman, and Sims (1984), and Meyer and Zaman (2013). Giannone, et al. (2014) use VARs to construct forecasts of inflation conditional on paths for oil and other price indicators. Baumeister and Kilian (2013) consider forecasts of oil prices conditioned on a range of scenarios. Schorfheide and Song (2013) and Aastveit, et al. (2014) use VARs to produce multi-step forecasts of growth, inflation, and other macroeconomic variables conditional on current-quarter forecasts obtained from other, judgmental sources (the Federal Reserve's Greenbook for the former and Survey of Professional Forecasters for the latter).

In light of common need for conditional forecasts, the attention paid to them, and their use in conveying the future stance of monetary policy, one would like to have a feel for their quality. Accordingly, in this paper, we develop and apply methods for the evaluation of conditional forecasts from VARs, using tests of bias, efficiency, and the MSE accuracy of conditional versus unconditional forecasts. More specifically, we provide analytical, Monte Carlo, and empirical evidence on tests of predictive ability for conditional forecasts from estimated models. In the empirical analysis, we consider forecasts of growth, unemployment, and inflation from a VAR, based on conditions on the short-term interest rate.

Throughout, our intention is to provide usable metrics for evaluating conditional forecasts, in a general sense and in comparison to the accuracy of unconditional forecasts. To do so, we focus on particular forms of conditional forecasts for which interpretation of various null and alternative hypotheses is most straight-forward. In particular, in our analysis, we consider forecasts conditioned on actual future information on some variables in the VAR model. In practice, conditional forecasts are sometimes constructed with future information (e.g., based on forward guidance from the central bank about policy rates), but not always. In general, though, the efficacy of conditional forecasts rests on having a properly

specified model. Our testing based on future information provides a way of assessing proper specification of the VAR. As we detail below, Herbst and Schorfheide (2012) use a similar idea in a Bayesian evaluation framework.

To better understand our approach to inference, consider a very simple example of a conditional forecast in which we forecast inflation (y_t) conditioned on a path for the federal funds rate (x_t) over the next two periods $t + 1$ and $t + 2$. Suppose that the assumed data-generating process for inflation and the funds rate is a zero-mean stationary VAR(1) taking the form

$$\begin{pmatrix} y_t \\ x_t \end{pmatrix} = \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \begin{pmatrix} y_{t-1} \\ x_{t-1} \end{pmatrix} + \begin{pmatrix} e_t \\ v_t \end{pmatrix},$$

with i.i.d. $N(0,1)$ errors with contemporaneous correlation ρ . Following the approach taken in Doan, Litterman, and Sims (1984), conditional on this path for the funds rate, the minimum mean square error (MSE) one- and two-step ahead forecasts of y_t are as follows:

$$\begin{aligned} \hat{y}_{t,1}^c &= \hat{y}_{t,1}^u + \hat{\rho}(\hat{x}_{t,1}^c - \hat{x}_{t,1}^u) \\ \hat{y}_{t,2}^c &= \hat{y}_{t,2}^u + (\hat{b} + \hat{\rho}(\hat{a} - \hat{c}))(\hat{x}_{t,1}^c - \hat{x}_{t,1}^u) + \hat{\rho}(\hat{x}_{t,2}^c - \hat{x}_{t,2}^u), \end{aligned}$$

where the superscripts c and u denote conditional and unconditional forecasts, respectively. In both cases the conditional forecasts of y are comprised of the standard, unconditional *MSE*-optimal forecast $\hat{y}_{t,j}^u$, $j = 1, 2$, plus additional terms that capture the impact of conditioning on future values of the federal funds rate, $\hat{x}_{t,1}^c$ and $\hat{x}_{t,2}^c$.

After rearranging terms, the conditional forecast errors $\hat{\varepsilon}_{t,2}^c = y_{t+2} - \hat{y}_{t,2}^c$ and $\hat{\varepsilon}_{t,1}^c = y_{t+1} - \hat{y}_{t,1}^c$ take the form

$$\begin{aligned} \hat{\varepsilon}_{t,1}^c &= \hat{\varepsilon}_{t,2}^u - \hat{\rho}(\hat{v}_{t,1}^u - \hat{v}_{t,1}^c) \\ \hat{\varepsilon}_{t,2}^c &= \hat{\varepsilon}_{t,2}^u - (\hat{b} + \hat{\rho}(\hat{a} - \hat{c}))(\hat{v}_{t,1}^u - \hat{v}_{t,1}^c) - \hat{\rho}(\hat{v}_{t,2}^u - \hat{v}_{t,2}^c). \end{aligned}$$

We immediately see that any “good” properties that the conditional forecast errors $\hat{\varepsilon}_{t,1}^c$ and $\hat{\varepsilon}_{t,2}^c$ have, such as unbiasedness or efficiency, are jointly determined by: (i) the quality of the unconditional forecast errors $\hat{\varepsilon}_{t,2}^u$, $\hat{\varepsilon}_{t,1}^u$, $\hat{v}_{t,1}^u$, and $\hat{v}_{t,2}^u$, as well as (ii) the behavior of the conditioning as measured via $\hat{v}_{t,1}^c$ and $\hat{v}_{t,2}^c$. As such, any method of inference designed to evaluate the quality of the conditional forecasts must somehow distinguish between properties determined by the quality of the model and properties determined by the quality of the conditioning.

Since these properties seem impossible to separate in general, we consider a simpler approach in which we condition on the ex-post realized values of the conditioning variables

so that $\hat{x}_{t,1}^c = x_{t+1}$ and $\hat{x}_{t,2}^c = x_{t+2}$. Since both $\hat{v}_{t,1}^c$ and $\hat{v}_{t,2}^c$ are then numerically zero we find that any properties associated with the conditional forecast errors $\hat{\varepsilon}_{t,1}^c$ and $\hat{\varepsilon}_{t,2}^c$ are determined only by the quality of the VAR as measured through the unconditional forecast errors, the regression parameters, and the residual variance parameters. Because we take this approach to conditioning, our inferential procedures are designed to evaluate the ability of the VAR to construct good conditional forecasts rather than evaluating a specific set of conditional forecasts per se.

While we focus on forecasts from time series models, conditional forecasts are routine in professional forecasting. The forecasts from the Federal Reserve’s Open Market Committee that have been published since 1979 are produced conditional on “appropriate monetary policy” as viewed by the respective individual members of the FOMC. In effect, the individual member of the FOMC is asked to produce a point forecast of, say, inflation over the next year given that the federal funds rate takes values over the coming year that are appropriate in the eyes of that individual FOMC member. Similarly, staff forecasts from the Federal Reserve Board are typically conditional on a future path of the funds rate as well as other variables.¹ The Bank of England, Riksbank, Norges Bank, and ECB all produce and release forecasts to the public that are conditional on a hypothetical future stance of monetary policy in one way or another.

Despite being explicitly conditional, the forecasts released by the FOMC and policymakers at other central banks seem to be regularly analyzed by the public and financial markets without taking account of the conditional nature of the forecasts. The same can also be said for academics. For example, Romer and Romer (2008) use standard MSE-based encompassing regressions to infer that Greenbook forecasts have an informational advantage over the corresponding FOMC forecasts. Patton and Timmermann (2012) apply newly developed tests of MSE-based forecast optimality to Greenbook forecasts and find evidence against forecast rationality. Gavin (2003), Gavin and Mandel (2003), Gavin and Pande (2008), Joutz and Stekler (2000), and Reifschneider and Tulip (2007) each evaluate the quality of FOMC forecasts in the context of MSEs.

However, two previous studies have developed methods for the evaluation of some form of conditional forecasts. Motivated by an interest in evaluating the efficiency of Greenbook forecasts, Faust and Wright (2008) develop a regression-based test of predictive ability

¹These forecasts are often referenced as Greenbook forecasts.

that accounts for the conditioning of the Greenbook forecasts on a pre-specified path of the federal funds rate over the forecast horizon. For the purpose of evaluating forecasts from DSGE models, Herbst and Schorfheide (2012) develop Bayesian methods to check the accuracy of point and density forecasts. More specifically, Herbst and Schorfheide consider the Bayesian tool of posterior predictive checks and forecasts of each variable conditioned on the actual future path of another, selected variable. Our paper differs from these in that we focus on conditional forecasts from VARs and emphasize frequentist inference.

The remainder of the paper proceeds as follows. Section 2 describes the two different approaches to conditional forecasting that we consider. Section 3 provides our theoretical results (proofs are provided in Appendix 1). Section 4 presents Monte Carlo evidence on the finite-sample accuracy of our proposed methods for evaluating conditional forecasts from VARs. Section 5 presents a practical application, to macroeconomic forecasts conditioned on interest rate paths. Section 6 concludes.

2 Conditional Forecasting Approaches

In generating and evaluating conditional forecasts, we consider two approaches that seem to be common in VAR and DSGE forecasting. All but one of our theoretical results apply to both approaches.

The standard in VAR forecasting is based on the textbook problem of conditional projection, as could be handled with a state space formulation of the VAR and the Kalman filter and smoother (see, e.g., Clarida and Coyle (1984) or Giannone, et al (2014)). The conditions on the variables of interest are contained in the measurement vector and equation; the data vector of the VAR is the state vector of the transition equation. The projection problem is one of predicting the state vector given the measurements (conditions). Doan, Litterman, and Sims (1984) developed an alternative approach to solving this formulation of the conditional forecasting problem, which consists of solving a least squares problem to pick the shocks needed to satisfy the conditions. In the context of conditioning on a policy path, this approach to conditional forecasting can be seen as consisting of the following: determining the set of shocks to the VAR that, by a least squares metric, best meet the conditions on the policy rate. In practice, this approach may mean that, in a given episode, an unchanged path of the policy rate could, for example, be due to shocks to output. Under this approach, the conditional forecasts are **not** dependent on the identification of struc-

tural shocks in the VAR. Note that under this approach, the forecast for each period can be affected by the imposition of conditions at all periods. For example, if we impose conditions for two periods, the forecast for period $t + 2$ will generally be affected by the conditions on both period $t + 1$ and $t + 2$.

In the interest of simplicity, in our implementation of minimum-MSE form forecasting, we abstract from the enhancements developed in Waggoner and Zha (1999).² In our implementation, as is typical, we estimate VAR parameters without taking the forecast conditions into account. Waggoner and Zha develop a Gibbs sampling algorithm that provides the exact finite-sample distribution of the conditional forecasts, by taking the conditions into account when sampling the VAR coefficients. Our reasons for abstracting from their extension are primarily computational. With the size of the large model we use, their Gibbs sampling algorithm would be extremely slow, due to computations of an extremely large VAR coefficient variance-covariance matrix. Moreover, based on a check we performed with a small BVAR model, their approach to conditional forecasting didn't seem to affect the conditional forecasts much.

In DSGE forecasting, the more standard approach for achieving conditions on the policy path rests on feeding in structural shocks to monetary policy needed to hit the policy path (see, e.g., Del Negro and Schorfheide (2013)). We use the following approach to implementing such an approach with our VARs. Under this approach, the scheme for identifying policy shocks does matter for the conditional forecasts. With our focus on conditioning on the policy path, it is the identification of monetary policy that matters. Accordingly, we refer to the forecasts we obtain under this approach as *conditional-policy shock* forecasts. Following common precedent with models such as ours (including, e.g., Bernanke and Blinder (1992), Christiano, Eichenbaum, and Evans (1996), and Giannone, Lenza, and Primiceri (2012)), we rely on identification of monetary policy via a recursive ordering of model innovations. Conditional forecasting under this approach involves the steps listed below. Note that, under this approach, as applied with a VAR, the forecast in period $t + \tau$ is not affected by conditions imposed on future periods $t + \tau + 1$, etc.

1. Using a model estimated with data through period t , form 1-step ahead forecasts for period $t + 1$.
2. Compute the structural shock to the federal funds rate needed to make the federal

²Jarocinski (2010) simplifies the formula of Waggoner and Zha (1999) and develops an extension.

funds rate equal the conditioning rate, and store that value in a vector of (VAR) innovations $\tilde{\varepsilon}_t$, where $\tilde{\varepsilon}_t$ contains 0 in all positions except that associated with the federal funds rate. Using S = the Choleski factor of the variance-covariance matrix of structural shocks, compute the implied reduced-form shocks as $S\tilde{\varepsilon}_t$. Re-compute 1-step ahead forecasts using this vector of shocks.

3. Move forward a step, and repeat steps 1-2 using the preceding period's forecast as data for the preceding period.
4. Continue through period $t + \tau$ to obtain forecasts from $t + 1$ through $t + \tau$.

3 Analytical results

We present our theoretical results in an environment in which OLS-estimated VARs are used to construct τ -step ahead conditional forecasts sequentially across forecast origins $t = R, \dots, T - \tau = R + P - \tau$. Specifically, suppose that the model takes the form

$$Y_t = C + A(L)Y_{t-1} + \varepsilon_t,$$

where $Y = (y_1, y_2, \dots, y_n)'$, $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n)'$, and $A(L) = \sum_{j=1}^l A_j L^j$ for $n \times 1$ and $n \times n$ parameter matrices C and A_j , $j = 1, \dots, l$, respectively. This is equivalent to

$$Y_t = \Lambda x_{t-1} + \varepsilon_t = (x'_{t-1} \otimes I_n)\beta + \varepsilon_t$$

if we define $x_t = (1, Y'_t, \dots, Y'_{t-l+1})'$, $\beta = \text{vec}(\Lambda)$, and $\Lambda = (C, A_1, \dots, A_l)$. If the model is estimated using the recursive scheme we obtain the estimated regression parameters

$$\hat{\beta}_t = \text{vec}(\hat{\Lambda}_t) = \text{vec}\left((t^{-1} \sum_{s=1}^{t-1} Y_{s+1} x'_s)(t^{-1} \sum_{s=1}^{t-1} x_s x'_s)^{-1}\right)$$

and corresponding residual variance matrix

$$\hat{\Sigma}_t = t^{-1} \sum_{s=1}^{t-1} \hat{\varepsilon}_{s+1} \hat{\varepsilon}'_{s+1}.$$

If the model is estimated using a rolling window of R observations we obtain comparable estimators but defined over the observations $s = t - R + 1, \dots, t$ rather than $s = 1, \dots, t$.

Both the regression and variance estimates $\hat{\beta}_t$ and $\hat{\Sigma}_t$ are used to construct iterated multi-step unconditional and conditional forecasts of Y_{t+j} , denoted $\hat{Y}_{t,j}^u$ and $\hat{Y}_{t,j}^c$, which in turn imply forecast errors $\hat{\varepsilon}_{t,j}^u$ and $\hat{\varepsilon}_{t,j}^c$. If we are interested in forecasting the i th element of $Y_{t+\tau}$, define $\hat{y}_{i,t,\tau} = \iota_i' \hat{Y}_{t,\tau}$ for the vector ι_i with a 1 in the i th position and zeroes elsewhere.

Define $\hat{\varepsilon}_{i,t,j}^c = \iota_i' \hat{\varepsilon}_{t,j}^c = y_{i,t+j} - \hat{y}_{i,t,j}^c$ and $\hat{\varepsilon}_{i,t,j}^u = \iota_i' \hat{\varepsilon}_{t,j}^u = y_{i,t+j} - \hat{y}_{i,t,j}^u$, accordingly. To simplify notation, and without loss of generality, we assume that the first element of Y is the primary element of interest and hence most of our analysis emphasizes the properties of the conditional forecast $\hat{y}_{1,t,j}^c$ and associated forecast error $\hat{\varepsilon}_{1,t,j}^c$.

As discussed in Section 2, the conditional forecasts are assumed to be either minimum-MSE or policy shock-based. In both cases it can be shown that the forecasts take the form

$$\hat{y}_{1,t,\tau}^c = \hat{y}_{1,t,\tau}^u + \sum_{i=1}^n \sum_{j=1}^m \hat{\gamma}_{i,t,j} (\hat{y}_{i,t,j}^u - \hat{y}_{i,t,j}^c)$$

for a collection of constants $\hat{\gamma}_{i,j}$ that are non-stochastic functions of both $\hat{\beta}_t$ and $\hat{\Sigma}_t$.³ Note that this structure aligns with the simple example from the introduction. In addition, we allow the maximum conditioning horizon m to be greater than or less than the forecast horizon τ , and we allow for direct conditioning on some elements of the future path of y_1 itself.

As mentioned earlier, we focus on evaluating the ability of the model to construct good conditional forecasts rather than evaluating conditional forecasts per se. We do so by examining the properties of the conditional forecast error when we condition on future realized values of those variables in the hypothetical scenario of interest. Specifically, if we set $\hat{y}_{i,t,j}^c = y_{i,t+j}$ we obtain

$$\hat{\varepsilon}_{1,t,\tau}^c = \hat{\varepsilon}_{1,t,\tau}^u - \sum_{i=1}^n \sum_{j=1}^m \hat{\gamma}_{i,t,j} \hat{\varepsilon}_{i,t,j}^u.$$

By taking this approach, the conditional forecast error has a representation as a linear function of unconditional forecast errors across all variables in the scenario and across all conditioning horizons.

Before proceeding to the tests we need to introduce some notation. Define $\hat{\phi}_t = (\hat{\beta}_t', \text{vech}(\hat{\Sigma}_t)')'$, $\phi = (\beta', \text{vech}(\Sigma)')'$, $B_\beta = ((Ex_s x_s')^{-1} \otimes I_n)$, $B_\Sigma = I_{n(n+1)/2}$, $B = \text{diag}(B_\beta, B_\Sigma)$, and $h_{s+1} = (h'_{\beta,s+1}, h'_{\Sigma,s+1})' = (\text{vec}(\varepsilon_{s+1} x_s')', \text{vech}(\varepsilon_{s+1} \varepsilon_{s+1}' - \Sigma)')'$. For any parametric function $z_t(\cdot)$, let $\hat{z}_t = z_t(\hat{\phi}_t)$ and $z_t = z_t(\phi)$. Finally, the asymptotics we use require both the initial in-sample size R and number of forecasts P to diverge as the overall sample increases. By taking this approach we find that the asymptotic distribution is influenced by the ratio $\lim_{P,R \rightarrow \infty} P/R = \pi$. In particular, the asymptotic variances developed below

³As a practical matter, many of the $\gamma_{i,j}$ will be zero depending on how many variables are conditioned on and how long the maximal conditioning horizon is. In the assumptions, we impose the restriction that $\gamma_{1,\tau}$ is zero (and hence the value of $\hat{y}_{1,t,\tau}^c$ is not imposed directly) via rank conditions on certain variance matrices.

all depend on the weights λ_{fh} and λ_{hh} as described in the following table taken from West and McCracken (1998).

	$\lambda_{fh} =$	$\lambda_{hh} =$
Recursive	$1 - \pi^{-1} \ln(1 + \pi)$	$2(1 - \pi^{-1} \ln(1 + \pi))$
Rolling, $\pi \leq 1$	$\pi/2$	$\pi - \pi^2/3$
Rolling, $1 < \pi < \infty$	$1 - (2\pi)^{-1}$	$1 - (3\pi)^{-1}$

3.1 Regression based tests of bias and efficiency

In this section we develop tests of zero bias and efficiency in the context of conditional forecasts from VARs when conditioning on future values of the variables in a hypothetical scenario. To do so first note that each can be couched in the context of a test of the null hypothesis that the coefficient α is zero in the regression

$$\hat{\varepsilon}_{1,t,\tau}^c = \hat{g}_t' \alpha + error, \quad t = R, \dots, T - \tau$$

for appropriate definitions of \hat{g}_t . Examples include a test of zero bias if we let $\hat{g}_t = 1$ and a test of efficiency if we let $\hat{g}_t = (1, \hat{y}_{1,t,\tau}^u)$ or perhaps $\hat{g}_t = (1, \hat{y}_{1,t,\tau}^u, \hat{y}_{i,t,j}^u)$ for those variables y_i in the scenario conditioned at horizon j .⁴

In each case, $P^{1/2} \hat{\alpha}$ is asymptotically normal with zero mean and a variance that accounts for estimation error in the estimated conditional forecast errors $\hat{\varepsilon}_{1,t,\tau}^c$ and generated regressors \hat{g}_t . These results follow directly from Lemmas 4.1 - 4.3 and Theorems 4.1 and 4.2 of West and McCracken (1998) if, under the null that $\alpha = 0$, we maintain a correctly specified VAR with errors ε_t that form a martingale difference sequence. Nevertheless it is worth noting that the *interpretation* of the results is slightly different than that intended in West and McCracken (1998). This arises because the conditional forecast errors depend on $\hat{\phi}_t = (\hat{\beta}_t', \text{vech}(\hat{\Sigma}_t)')$ rather than just $\hat{\beta}_t$. Nevertheless, $\hat{\phi}_t$ satisfies assumption A2 of West and McCracken (1998) and the remainder of their assumptions are satisfied except for A1 (c) and (d).⁵ These assumptions, however, do not affect the derivation of asymptotic normality. They are only used in Theorem 5.1 of West and McCracken (1998), wherein special cases are delineated under which estimation error is asymptotically irrelevant. For clarity we restate the assumptions in the context of the current paper and then proceed to the Theorem.

⁴Note that this last regression is not the standard Mincer-Zarnowitz form of the efficiency regression. We consider this separately in the next section.

⁵See section 3.4.3 of Lutkepohl (1991) for details.

Assumption 1: (a) In some neighborhood N around ϕ , and with probability 1, $\varepsilon_{1,t,\tau}^c(\phi)$ and $g_t(\phi)$ are measurable and twice continuously differentiable; (b) $E(\varepsilon_t|x_{t-j}, \varepsilon_{t-j} \text{ all } j \geq 1) = 0$; (c) $Eg_t g_t'$ is full rank.

Assumption 2: The estimate $\hat{\phi}_t$ satisfies $\hat{\phi}_t - \phi = B(t)H(t)$, with (a) $B(t) \rightarrow_{a.s.} B = \text{diag}[(Ex_s x_s')^{-1} \otimes I_n, I_{n(n+1)/2}]$ and B of full rank; (b) $H(t)$ equals $t^{-1} \sum_{s=1}^{t-1} h_{s+1}$ and $t^{-1} \sum_{s=t-R+1}^{t-1} h_{s+1}$ for the recursive and rolling schemes, respectively; (c) $Eh_{s+1} = 0$; (d) In the neighborhood N of Assumption 1, h_{s+1} is measurable and continuously differentiable.

Assumption 3: In the neighborhood N of Assumption 1, there is a constant $D < \infty$ such that for all t , $\sup_{\phi \in N} |\partial^2 \varepsilon_{1,t,\tau}^c(\phi) / \partial \phi \partial \phi'| < m_t$ for a measurable m_t for which $Em_t^4 < D$. The same holds when $\varepsilon_{1,t,\tau}^c(\phi)$ is replaced by $g_t(\phi)$.

Assumption 4: Let $w_t = (x_t', \text{vec}(\partial g_t / \partial \phi)', \varepsilon_t', g_t', h_t')'$. (a) For some $d > 1$, $\sup_t E\|w_t\|^{8d} < \infty$, where $\|\cdot\|$ denotes the Euclidean norm; (b) w_t is strong mixing with coefficients of size $-3d/(d-1)$; (c) w_t is fourth-order stationary; (d) $S_{ff} = \lim_{P,R \rightarrow \infty} \text{Var}(P^{-1/2} \sum_{t=R}^{T-\tau} \varepsilon_{1,t,\tau}^c g_t)$ is positive definite.

Assumption 5: $R, P \rightarrow \infty$ as $T \rightarrow \infty$ with $\lim_{T \rightarrow \infty} \frac{P}{R} = \pi$. (a) $0 \leq \pi \leq \infty$ for the recursive scheme; (b) $0 \leq \pi < \infty$ for the rolling scheme.

Theorem 1 *Maintain assumptions A1-A5. $P^{1/2} \hat{\alpha} \rightarrow^d N(0, V)$ with $V = (Eg_t g_t')^{-1} \Omega (Eg_t g_t')^{-1}$ and*

$$\Omega = S_{ff} + \lambda_{fh}(FBS_{fh} + S'_{fh}B'F') + \lambda_{hh}FBS_{hh}B'F',$$

where $S_{ff} = \lim_{P,R \rightarrow \infty} \text{Var}(P^{-1/2} \sum_{t=R}^{T-\tau} \varepsilon_{1,t,\tau}^c g_t)$, $S_{hh} = \lim_{T \rightarrow \infty} \text{Var}(T^{-1/2} \sum_{s=1}^{T-1} h_{s+1})$, $S_{fh} = \lim_{P,R \rightarrow \infty} \text{Cov}(P^{-1/2} \sum_{t=R}^{T-\tau} \varepsilon_{1,t,\tau}^c g_t, P^{-1/2} \sum_{t=R}^{T-\tau} h'_{s+1})$, and $F = E(\partial \varepsilon_{1,t,\tau}^c(\phi) g_t(\phi) / \partial \phi)$.

As in West and McCracken (1998), the asymptotic variance, especially through Ω , is comprised of three components: S_{ff} captures the variation that would exist even if ϕ were known, $FBS_{hh}B'F'$ captures the variation due purely to estimation error in $\hat{\phi}_t$, and $FBS_{fh} + S'_{fh}B'F'$ is the covariance between the two sources of variability.

The asymptotic variance is complicated but can be simplified for each of the tests discussed above. First, note that since $E\varepsilon_{1,t,\tau}^c(\partial g_t(\phi) / \partial \phi) = 0$, F trivially reduces to $Eg_t(\partial \varepsilon_{1,t,\tau}^c(\phi) / \partial \phi)$. In addition, it is straightforward to show that $Eg_t(\partial \varepsilon_{1,t,\tau}^c(\phi) / \partial \text{vech}(\Sigma)) = 0$ and hence $F = (Eg_t(\partial \varepsilon_{1,t,\tau}^c(\phi) / \partial \beta), 0)$. That, along with the fact that B is block diagonal, implies that in large samples there is no estimation error contributed by $\hat{\Sigma}_t$ in either

\hat{g}_t or $\hat{\varepsilon}_{1,t,\tau}^c$. Notationally, we can therefore simplify the formula for Ω to

$$\Omega = S_{ff} + \lambda_{fh}(F_\beta B_\beta S_{fh,1} + S'_{fh,1} B'_\beta F'_\beta) + \lambda_{hh} F_\beta B_\beta S_{hh,11} B'_\beta F'_\beta,$$

where $F_\beta = E g_t(\partial \varepsilon_{1,t,\tau}^c(\phi)/\partial \beta)$ and $S_{fh,1}$ and $S_{hh,11}$ denote those elements of S_{fh} and S_{hh} associated only with the OLS moment conditions $h_{\beta,s+1}$.

With a bit more work, F can be explicitly derived using elements of section 3.5.2 of Lutkepohl (1991). Specifically if we define the $(nl + 1) \times (nl + 1)$ matrix

$$W = \begin{pmatrix} 1 & 0 & \dots & 0 \\ & \Lambda & & \\ 0 & I_{n(l-1)} & & 0 \end{pmatrix},$$

the $n \times (nl + 1)$ selection matrix $J_1 = (0_{n \times 1}, I_n, 0_{n \times n(l-1)})$, $\Phi_i = J_1 W^i J'_1$, and $\Theta_{i,t,j} = \iota'_i \sum_{q=0}^{j-1} (x'_t(W')^{j-1-q} \otimes \Phi_q)$, we obtain

$$F_\beta = E g_t(-\Theta_{1,t,\tau} + \sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j} \Theta_{i,t,j}).$$

3.2 Mincer-Zarnowitz efficiency test

In the previous section we delineated a regression-based test of efficiency based on a regression of $\hat{\varepsilon}_{1,t,\tau}^c$ on $(1, \hat{y}_{1,t+\tau}^u)$. By taking this approach we were able to use the results in West and McCracken (1998) to prove that the regression coefficients are asymptotically normal under the null hypothesis that $\alpha = 0$. While useful, this result deviates from a more standard version of the Mincer and Zarnowitz (1969) test of efficiency in which, for conditional forecasts, we would regress $\hat{\varepsilon}_{1,t,\tau}^c$ on $(1, \hat{y}_{1,t,\tau}^c)$.

The reason we didn't consider the Mincer-Zarnowitz regression is that, by conditioning on future values of the variables in the scenario, it is possible that $E \hat{\varepsilon}_{1,t,\tau}^c \hat{y}_{1,t+\tau}^c \neq 0$ even though $\hat{\varepsilon}_{1,t,\tau}^c$ is orthogonal to all information in the time t information set. As it turns out, whether or not that orthogonality restriction holds depends on whether minimum-MSE or policy shock-based conditioning is used. Straightforward algebra show that, under minimum-MSE, but not policy shock, conditioning, $E \hat{\varepsilon}_{1,t,\tau}^c \hat{y}_{1,t+\tau}^c = 0$ and hence Theorem 1 is indeed applicable with $\hat{g}_t = (1, \hat{y}_{1,t,\tau}^c)'$. Even so, the previously discussed simplifications of the asymptotic variance Ω no longer hold. In particular, under minimum-MSE conditioning, $\hat{\Sigma}_t$ contributes estimation error even in the limit. We can see this directly in the following derivation of the $(2 \times (\dim \text{vec}(\Lambda) + \dim \text{vech}(\Sigma)))$ matrix $F = (F'_1, F'_2)' =$

$E(\partial\varepsilon_{1,t,\tau}^c(\phi)g_t(\phi)/\partial\phi)$:

$$\begin{aligned}
F_1 &= (E(-\Theta_{1,t,\tau} + \sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j}\Theta_{i,t,j}), 0_{n(n+1)/2}) \\
F_2 &= (E(y_{1,t,\tau}^u)(-\Theta_{1,t,\tau} + \sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j}\Theta_{i,t,j}) - E(y_{1,t,\tau}^c - y_{1,t,\tau}^u)(\sum_{i=1}^n \sum_{j=1}^m \nabla_{\beta}\gamma_{i,j}u_{i,t,j}^u) \\
&\quad + E(u_{1,t,\tau}^c)(\sum_{i=1}^n \sum_{j=1}^m \nabla_{\beta}\gamma_{i,j}u_{i,t,j}^u), -E(y_{1,t,\tau}^c)(\sum_{i=1}^n \sum_{j=1}^m \nabla_{\Sigma}\gamma_{i,j}u_{i,t,j}^u) \\
&\quad + E(u_{1,t,\tau}^c)(\sum_{i=1}^n \sum_{j=1}^m \nabla_{\Sigma}\gamma_{i,j}u_{i,t,j}^u)).
\end{aligned}$$

The first row of F is identical to that for the test of zero bias (as expected), concatenated by zeros, since $E\partial\varepsilon_{1,t,\tau}^c(\phi)/\partial\text{vech}(\Sigma) = 0$. In contrast, $E\partial\varepsilon_{1,t,\tau}^c(\phi)y_{1,t,\tau}^c(\phi)/\partial\text{vech}(\Sigma) \neq 0$, and hence the second row of F includes terms contributing to the asymptotic variance due to estimation error in $\hat{\Sigma}_t$.

While useful under minimum-MSE conditioning, we are nevertheless left with the fact that the Mincer-Zarnowitz regression is not applicable under policy shock-based conditioning. A related point is made by Faust and Wright (2008) in the context of testing the efficiency of the Greenbook forecasts constructed by the staff at the Federal Reserve Board of Governors. To get around this problem they suggest an alternative formulation of the Mincer-Zarnowitz regression designed to “soak up” the impact of conditioning on the values of variables not in the time t information set. In the notation of our paper, as well as our approach to conditioning, their regression takes the form

$$\begin{aligned}
\hat{\varepsilon}_{1,t,\tau}^c &= \alpha_0 + \alpha_1(\hat{y}_{1,t,\tau}^u - \hat{y}_{1,t,\tau}^c) + \alpha_2\hat{y}_{1,t,\tau}^c + \text{error} \\
&= \hat{g}_t'\alpha + \text{error},
\end{aligned} \tag{1}$$

where $\hat{g}_t = (1, \hat{y}_{1,t,\tau}^u - \hat{y}_{1,t,\tau}^c, \hat{y}_{1,t,\tau}^c)'$.

At first blush this regression looks like those discussed in the previous section. And yet there is one major difference that precludes directly applying the results in West and McCracken (1998): α_1 need not be zero under the null of efficiency which, here, is represented by $\alpha_2 = 0$ (and perhaps $\alpha_0 = 0$ if a joint test of efficiency and zero bias is desired). Since the proofs in West and McCracken (1998) explicitly require that $E\hat{\varepsilon}_{1,t,\tau}^c\hat{g}_t = 0$, and hence all elements of α are zero, their results are not applicable.

Regardless, asymptotic normality of the coefficients can be established using the more general results in West (1996) along with the Delta method. To do so, first define the (8×1) function $f_{t,\tau}(\hat{\phi}_t) = (\text{vech}_{-1}(\hat{g}_t\hat{g}_t'), \hat{g}_t'\hat{\varepsilon}_{1,t,\tau}^c)'$, where the notation vech_{-1} denotes the vech operator but omits the first element (that associated with the intercept). There then exists a

twice continuously differentiable function $q(\cdot): \mathbb{R}^8 \rightarrow \mathbb{R}^3$ satisfying $\hat{\alpha} = q(P^{-1} \sum_{t=R}^{T-\tau} f_{t,\tau}(\hat{\phi}_t))$ such that $\alpha_0 = \iota_1' q(Ef_{t,\tau}) = 0$ and $\alpha_2 = \iota_3' q(Ef_{t,\tau}) = 0$ under the null hypothesis. Finally, define $\nabla q(Ef_{t,\tau}) = \partial q(Ef_{t,\tau}) / \partial Ef_{t,\tau}$. The following Theorem provides the asymptotic distribution of $P^{1/2}(\hat{\alpha} - \alpha)$.

Theorem 2 *Maintain assumptions A1-A5. $P^{1/2}(\hat{\alpha} - \alpha) \rightarrow^d N(0, V)$ with $V = \nabla q(Ef_{t,\tau})' \Omega \nabla q(Ef_{t,\tau})$ and*

$$\Omega = S_{ff} + \lambda_{fh}(FBS_{fh} + S'_{fh}B'F') + \lambda_{hh}FBS_{hh}B'F',$$

where $S_{ff} = \lim_{P,R \rightarrow \infty} \text{Var}(P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau})$, $S_{fh} = \lim_{P,R \rightarrow \infty} \text{Cov}(P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}, P^{-1/2} \sum_{t=R}^{T-\tau} h'_{s+1})$, $S_{hh} = \lim_{T \rightarrow \infty} \text{Var}(T^{-1/2} \sum_{s=1}^{T-1} h_{s+1})$, and $F = E(\partial f_{t,\tau}(\phi) / \partial \phi)$.

Again we find that the regression coefficients are asymptotically normal. But in contrast to the results in the previous section, only α_0 and α_2 are necessarily zero when the VAR is correctly specified.

Also, in contrast to the previous results, there does not appear to be any way of simplifying the formulation of the asymptotic variance V . Both F_β and F_Σ are non-zero and hence estimation error from both $\hat{\beta}_t$ and $\hat{\Sigma}_t$ contribute to the asymptotic variance. In particular, while tedious, the formulas for $F = (F'_1, \dots, F'_8)'$ ($8 \times (\dim \text{vec}(\Lambda) + \dim \text{vech}(\Sigma))$) can be derived explicitly and take the following form:

$$\begin{aligned} F_1 &= (E \sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j} \Theta_{i,t,j}, 0_{n(n+1)/2}) \\ F_2 &= (E(\Theta_{1,t,\tau} - \sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j} \Theta_{i,t,j}), 0_{n(n+1)/2}) \\ F_3 &= (2E(y_{1,t,\tau}^c - y_{1,t,\tau}^u)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\beta \gamma_{i,j} u_{i,t,j}^u), 2E(y_{1,t,\tau}^c - y_{1,t,\tau}^u)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\Sigma \gamma_{i,j} u_{i,t,j}^u)) \\ F_4 &= (E(\sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j} y_{1,t,\tau}^u \Theta_{i,t,j}) - 2E(y_{1,t,\tau}^c - y_{1,t,\tau}^u)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\beta \gamma_{i,j} u_{i,t,j}^u), \\ &\quad -2E(y_{1,t,\tau}^c - y_{1,t,\tau}^u)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\Sigma \gamma_{i,j} u_{i,t,j}^u)) \\ F_5 &= (2E(y_{1,t,\tau}^c)(-\Theta_{1,t,\tau} + \sum_{i=1}^n \sum_{j=1}^m (-\gamma_{i,j} \Theta_{i,t,j} + \nabla_\beta \gamma_{i,j} u_{i,t,j}^u)), 2E(y_{1,t,\tau}^c)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\Sigma \gamma_{i,j} u_{i,t,j}^u)) \\ F_6 &= (E(-\Theta_{1,t,\tau} + \sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j} \Theta_{i,t,j}), 0_{n(n+1)/2}) \\ F_7 &= (-E \sum_{i=1}^n \sum_{j=1}^m \nabla_\beta \gamma_{i,j} u_{i,t,j}^u u_{1,t,\tau}^c + E \sum_{i=1}^n \sum_{j=1}^m \nabla_\beta \gamma_{i,j} u_{i,t,j}^u (y_{1,t,\tau}^c - y_{1,t,\tau}^u), \\ &\quad -E \sum_{i=1}^n \sum_{j=1}^m \nabla_\Sigma \gamma_{i,j} u_{i,t,j}^u u_{1,t,\tau}^c + E \sum_{i=1}^n \sum_{j=1}^m \nabla_\Sigma \gamma_{i,j} u_{i,t,j}^u (y_{1,t,\tau}^c - y_{1,t,\tau}^u)) \\ F_8 &= (E(y_{1,t,\tau}^u)(-\Theta_{1,t,\tau} + \sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j} \Theta_{i,t,j}) - E(y_{1,t,\tau}^c - y_{1,t,\tau}^u)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\beta \gamma_{i,j} u_{i,t,j}^u) \\ &\quad + E(u_{1,t,\tau}^c)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\beta \gamma_{i,j} u_{i,t,j}^u), -E(y_{1,t,\tau}^c)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\Sigma \gamma_{i,j} u_{i,t,j}^u) \\ &\quad + E(u_{1,t,\tau}^c)(\sum_{i=1}^n \sum_{j=1}^m \nabla_\Sigma \gamma_{i,j} u_{i,t,j}^u)). \end{aligned}$$

3.3 Equal accuracy test

In the previous two sections we described tests of predictive ability related to the properties of the conditional forecast error $\hat{\varepsilon}_{1,t,\tau}^c$. In each case, the null hypothesis relates to a property of the forecast error that suggests that the model is useful for constructing conditional forecasts. Rejecting the null hypothesis is indicative of a flaw in the model that might cause future conditional forecasts to be ill-behaved.

One property we have not discussed is the accuracy of the conditional forecast. Intuitively, one would expect that after conditioning on future values of those variables in the hypothetical scenario, the conditional forecast would be more accurate than an unconditional forecast that does not utilize any future information. In other words, if our conditional forecast is good we would expect $E(\varepsilon_{1,t,\tau}^c)^2 - E(\varepsilon_{1,t,\tau}^u)^2 < 0$. In fact, as also noted in Herbst and Schorfheide (2012), when the forecasts are of the minimum-MSE variety this should be the case.⁶

Unfortunately, this “good” property of the conditional forecast does not give us a workable null hypothesis under which to derive an asymptotic distribution and conduct inference. In fact, given our methodological approach of conditioning on future observations of those variables in a scenario, the null hypothesis $E(\varepsilon_{1,t,\tau}^c)^2 - E(\varepsilon_{1,t,\tau}^u)^2 = 0$ would require the model to be misspecified. Because of this, one approach to inference would be to derive the asymptotic distribution of the moment condition $P^{-1/2} \sum_{t=R}^{T-\tau} ((\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2)$ assuming a misspecified VAR and reject the null in the lower tail of this distribution. Unfortunately, rejecting the null in the lower tail does not imply that the model is “good” in any sense since the model could still be misspecified and yet $E(\varepsilon_{1,t,\tau}^c)^2 - E(\varepsilon_{1,t,\tau}^u)^2 < 0$.

We therefore take a different approach to inference, one that continues our strategy of maintaining a correctly specified VAR under the null. Again, under minimum-MSE conditioning, correct specification implies the existence of a non-negative constant k satisfying $E(\varepsilon_{1,t,\tau}^c)^2 - E(\varepsilon_{1,t,\tau}^u)^2 = k$. This constant depends on the VAR regression parameters A_j and residual variance Σ and can be derived explicitly as a function of ϕ . To do so, first define $\Psi_j \Sigma^{1/2}$ as the matrix of orthogonalized impulse responses after j periods and let

$$D = \begin{pmatrix} \Sigma^{1/2} & 0 & 0 & 0 \\ \Psi_1 \Sigma^{1/2} & \Sigma^{1/2} & 0 & 0 \\ \dots & \dots & \Sigma^{1/2} & 0 \\ \Psi_{\max(\tau,m)} \Sigma^{1/2} & \Psi_{\max(\tau,m)-1} \Sigma^{1/2} & \Psi_1 \Sigma^{1/2} & \Sigma^{1/2} \end{pmatrix}.$$

⁶This need not hold under policy shock-based conditioning. As such, we do not consider this approach within the section.

Now let \tilde{D} denote the matrix formed by those rows in D associated with a conditioning restriction.⁷ For example, if $n = 2$ and we condition on future values of the second element of the VAR at both the first and second horizons ($m = 2$), \tilde{D} consists of the (2×4) matrix formed by stacking the second and fourth rows of D . Straightforward algebra then implies $k(\phi) = \iota_1' D \tilde{D}' (\tilde{D} \tilde{D}')^{-1} \tilde{D} D' \iota_1$.

With this constant $k(\phi)$ in hand we consider testing for “equal accuracy” (below, we will just generally refer to this as testing MSE accuracy) using the appropriately re-centered Diebold and Mariano (1996) and West (1996)-type test of predictive ability:

$$P^{1/2} \hat{\alpha} = P^{-1/2} \sum_{t=R}^{T-\tau} ((\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k(\hat{\phi}_T)).$$

Theorem 3 *Maintain assumptions A1-A5. $P^{1/2} \hat{\alpha} \rightarrow^d N(0, \Omega)$ with*

$$\begin{aligned} \Omega = & S_{ff} + 2\lambda_{fh} FBS_{fh} - 2\frac{\pi}{1+\pi} KBS_{fh} - 2\frac{\pi}{1+\pi} FBS_{hh} B' K' \\ & + \lambda_{hh} FBS_{hh} B' F' + \frac{\pi}{1+\pi} KBS_{hh} B' K', \end{aligned}$$

where $S_{fh} = \lim_{P,R \rightarrow \infty} Cov(P^{-1/2} \sum_{t=R}^{T-\tau} ((\varepsilon_{1,t,\tau}^c)^2 - (\varepsilon_{1,t,\tau}^u)^2 - k), P^{-1/2} \sum_{t=R}^{T-\tau} h'_{s+1})$, $S_{ff} = \lim_{P,R \rightarrow \infty} Var(P^{-1/2} \sum_{t=R}^{T-\tau} ((\varepsilon_{1,t,\tau}^c)^2 - (\varepsilon_{1,t,\tau}^u)^2 - k))$, $S_{hh} = \lim_{T \rightarrow \infty} Var(T^{-1/2} \sum_{s=1}^{T-1} h_{s+1})$, $F = E(\partial((\varepsilon_{1,t,\tau}^c(\phi))^2 - (\varepsilon_{1,t,\tau}^u(\phi))^2) / \partial \phi)$, and $K = \partial k(\phi) / \partial \phi$.

Theorem 3 implies that an appropriately centered DM/W test of equal predictive ability is asymptotically normal. While this is precisely what is shown in DM/W, there is an important distinction: estimation error affects the asymptotic distribution. This contrasts with the results in West (1996). In that paper it is shown that when the forecasts associated with two non-nested OLS estimated linear models are evaluated under quadratic loss, $F = 0$ (and for that matter, $K = 0$) and hence estimation error is asymptotically irrelevant.

Mathematically, the key feature driving this difference is the presence of the weights $\gamma_{i,j}$ as well as the centering constant k , and, in particular the fact that they are functions of ϕ . Let $\nabla_{\beta} \gamma_{i,j} = \partial \gamma_{i,j}(\phi) / \partial \beta$ and $\nabla_{\Sigma} \gamma_{i,j} = \partial \gamma_{i,j}(\phi) / \partial vech(\Sigma)$. Straightforward algebra reveals that $F = (F_{\beta}, F_{\Sigma})$, where

$$F_{\beta} = - \sum_{i=1}^n \sum_{j=1}^m \nabla_{\beta} \gamma_{i,j} E u_{1,t,\tau}^u u_{i,t,j}^u + 2E(\sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j} u_{i,t,j}^u) (\sum_{i=1}^n \sum_{j=1}^m \nabla_{\beta} \gamma_{i,j} u_{i,t,j}^u)$$

and

$$F_{\Sigma} = - \sum_{i=1}^n \sum_{j=1}^m \nabla_{\Sigma} \gamma_{i,j} E u_{1,t,\tau}^u u_{i,t,j}^u + 2E(\sum_{i=1}^n \sum_{j=1}^m \gamma_{i,j} u_{i,t,j}^u) (\sum_{i=1}^n \sum_{j=1}^m \nabla_{\Sigma} \gamma_{i,j} u_{i,t,j}^u).$$

⁷This notation is taken directly from Jarocinski (2010) except we substitute D for R .

Since F_β and F_Σ (as well as unreported K_β and K_Σ) are both non-zero, estimation error in both $\hat{\beta}_t$ and $\hat{\Sigma}_t$ affect the asymptotic distribution through the asymptotic variance.

3.4 A bootstrap approach to inference

For each test of bias and efficiency we are able to establish that $P^{1/2}(\hat{\alpha} - \alpha)$ is asymptotically normal. We are also able to establish asymptotic normality of the test of MSE accuracy. As such, one approach to inference is to construct a consistent estimate of the asymptotic variance and then compare the standardized statistic $P^{1/2}(\hat{\alpha}_j - \alpha_j)/\hat{\Omega}_j^{1/2}$ to standard normal critical values. As discussed in West (1996) as well as West and McCracken (1998), many of the components of the asymptotic variance are easily estimated, while other elements can be more complicated. Among the easiest components are $\hat{\pi} = P/R$, $\hat{B}_\beta = ((T^{-1} \sum_{s=1}^{T-1} x_s x_s')^{-1} \otimes I_n)$, and $B_\Sigma = I_{n(n+1)/2}$. In addition, standard HAC estimators can be used to construct estimates of S_{ff} , S_{fh} , and S_{hh} given $f_{t,\tau}(\hat{\phi}_t)$ and $\hat{h}_{t+1} = (\text{vec}(\hat{\varepsilon}_{t+1} x_t')', \text{vech}(\hat{\varepsilon}_{t+1} \hat{\varepsilon}_{t+1}' - \hat{\Sigma}_T)')$, where $\hat{\varepsilon}_{t+1}$ denotes the residuals from the OLS-estimated VAR.

Unfortunately, estimating F , $\nabla q(Ef_{t,\tau})$, and K is significantly more difficult even when, for example, formulas for F are provided as they are in the text. As a result, we suggest and use in our applications a parametric bootstrap approach to inference. One might rely on a bootstrap patterned on the approach first used in Kilian (1999) and applied in the univariate forecast evaluation of such studies as Clark and McCracken (2001, 2005). Goncalves and Kilian (2004) establish the general asymptotic validity of such a bootstrap for many purposes. However, it is not necessarily valid for our purposes, because the asymptotic distributions of interest depend on the VAR's error variance matrix, Σ , a dependency outside the scope of the Goncalves-Kilian results. We instead use a VAR-based, moving-block residual based bootstrap developed in Bruggemann, Jentsch, and Trenkler (2014), which is valid in the presence of dependency of asymptotic distributions on Σ (as well as conditional heteroskedasticity of the VAR's innovations). To implement their bootstrap in our setting, we estimate the VAR of interest with the full sample of data and save the residuals and coefficients. We then bootstrap the residuals using a moving block bootstrap of 40 observations.⁸ We use these residuals and the autoregressive structure of the VAR to obtain an artificial time series y_t^* .⁹ In each bootstrap replication, the bootstrapped data are

⁸In our Monte Carlo analysis, before settling on a block size of 40, we considered a grid of sizes up through 50, and found differences across block choices to be fairly small.

⁹The initial observations are selected by sampling from the actual data as in Stine (1987).

used to estimate the VAR forecasting model at each forecast origin and generate artificial, out-of-sample forecasts. These forecasts and associated forecast errors are used to produce bias, efficiency, and MSE accuracy test statistics. Critical values are simply computed as percentiles of the bootstrapped test statistics.

3.5 Tests applied to conditional forecasts under Giacomini-White asymptotics

The theoretical results in West (1996) and West and McCracken (1998) focus on testing a null hypothesis of the form $H_0 : Ef_{t,\tau}(\phi) = 0$. In words, this hypothesis is designed to evaluate the predictive ability of a model if the true parameter values ϕ of the model are known. But one might prefer to evaluate the predictive content of the model accounting for the fact that the model parameters must be estimated and will never be known. Doing so changes the null hypothesis to something akin to $H_0: Ef_{t,\tau}(\hat{\phi}_t) = 0$, where the estimation error associated with the parameter estimates $\hat{\phi}_t$ is now captured under the null hypothesis.

Even though the null hypothesis $H_0: Ef_{t,\tau}(\hat{\phi}_t) = 0$ is reasonably intuitive, the results in West (1996) and West and McCracken (1998) do not apply under this null and hence their theory cannot be used to infer the asymptotic distribution of $P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}(\hat{\phi}_t)$ under this null. Instead, theoretical results in Giacomini and White (2006) can be used to conduct inference. Giacomini and White show that so long as all parameter estimates are estimated using a small rolling window of observations (small in the sense that R is finite and P diverges to infinity), $P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}(\hat{\phi}_t) \rightarrow^d N(0, S_{\hat{f}\hat{f}})$ under the null hypothesis

$$\lim_{P \rightarrow \infty} P^{-1/2} \sum_{t=R}^{T-\tau} Ef_{t,\tau}(\hat{\phi}_t) = 0,$$

where $S_{\hat{f}\hat{f}} = \lim_{P \rightarrow \infty} Var(P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}(\hat{\phi}_t))$.

The theoretical results in Giacomini and White (2006) can be used for a wide range of applications, including tests of zero bias and forecast efficiency applied to conditional forecasts.¹⁰ So long as the statistic takes the form $P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}(\hat{\phi}_t)$, and all parameters are estimated using a small rolling window of observations, normal critical values can be used to conduct inference on the null hypothesis $\lim_{P \rightarrow \infty} P^{-1/2} \sum_{t=R}^{T-\tau} Ef_{t,\tau}(\hat{\phi}_t) = 0$, without any additional correction of standard errors for the effects of estimation of the forecasting model's parameters. In practice, as much as it may be asymptotically valid (under a rolling estimation scheme for the forecasting model) to compare bias and efficiency tests against

¹⁰Because our test of equal MSE requires an estimated recentering term, and is estimated using the full sample, the Giacomini and White (2006) results are not directly applicable.

normal distributions without correction for parameter estimation error, HAC estimation problems may still make a bootstrap like the one we described above preferable to standard normal-based inference.

4 Monte Carlo Analysis

This section presents a Monte Carlo analysis of the finite-sample properties of tests for bias, efficiency, and MSE accuracy applied to unconditional and conditional forecasts from VAR models. In all cases, consistent with the preceding analytics, we produce and evaluate forecasts conditioned on a path for a pseudo-policy variable that is the actual future path. Using the simulated data, we conduct forecast inference based on both normal distributions with standard errors that abstract from parameter estimation error and bootstrapped test distributions.

In these experiments, we use bivariate and trivariate VARs as data-generating processes (DGPs). In the reported results, we form (iterated multi-step) forecasts using OLS estimates of VARs. In results not reported in the interest of brevity, we obtained very similar results for VARs estimated with Bayesian methods, under a Normal-inverted Wishart prior and posterior. We focus on forecasts computed under a recursive (expanding window) estimation scheme, but we provide some results for a rolling window estimation scheme. For the conditional forecasts, we concentrate on the minimum-MSE approach to conditional forecasting; we provide a briefer set of results for the policy shock approach to conditional forecasting.

In all simulations, based on 2000 Monte Carlo draws, we report the percentage of Monte Carlo trials in which the null of no bias, efficiency, or MSE accuracy is rejected — the percentage of trials in which the sample test statistics fall outside (two-sided) critical values. In the reported results, the tests are compared against 10% critical values. Using 5% critical values yields similar findings.

We proceed by first detailing the data-generating processes and other aspects of experiment design and then presenting the results.

4.1 Monte Carlo design

For each DGP, we generate data using draws of innovations and the autoregressive structure of the DGP. The initial observations necessitated by the lag structure of each DGP are

generated with draws from the unconditional normal distribution implied by the DGP. With quarterly data in mind, we report results for forecast horizons of 1, 2, and in some cases, 4 periods. We consider sample sizes of $R, P = 50,100; 50,150; 100,50; \text{ and } 100,100$. We use DGPs 1, 1G, and 2 to evaluate size properties and DGPs 3 and 4 to evaluate power.

DGP 1 is a bivariate VAR(1), with regression coefficients given in the first panel of Table 1 and an error variance-covariance matrix of:

$$\text{var}(\varepsilon_t) = \begin{pmatrix} 1.0 & \\ 0.5 & 1.0 \end{pmatrix}.$$

In order to assess test reliability with conditionally heteroskedastic forecast errors, we also consider a version of DGP 1 with GARCH innovations, denoted DGP 1G. The VAR has the same regression coefficients as given for DGP 1 in the first panel of Table 1. In the GARCH structure, taken from an experiment design in Bruggemann, Jentsch, and Trenkler (2014), let e_t denote a 2×1 innovation vector that is distributed $N(0, I_2)$. Let $v_{i,t} = \sigma_{i,t} e_{i,t}$, where $\sigma_{i,t}^2 = 0.2 + 0.05v_{i,t-1}^2 + 0.75\sigma_{i,t-1}^2$, such that $v_{1,t}$ and $v_{2,t}$ are independent GARCH(1,1) processes. The VAR innovations are constructed as Sv_t , where S is

$$\text{var}(\varepsilon_t) = \begin{pmatrix} 1.0 & 0.0 \\ 0.5 & 0.866 \end{pmatrix}.$$

With this formulation, the unconditional error variance matrix is the same as in the homoskedastic version of DGP 1.

DGP 2 is a trivariate VAR(2), with regression coefficients given in the second panel of Table 1 and an error variance-covariance matrix of:

$$\text{var}(\varepsilon_t) = \begin{pmatrix} 9.265 & & \\ 0.296 & 1.746 & \\ 0.553 & 0.184 & 0.752 \end{pmatrix}.$$

We set the parameters of DGP 2 to equal OLS estimates of a VAR in GDP growth, inflation less a survey-based measure of trend inflation (see, e.g., Clark and Doh (2014), Faust and Wright (2013)), and the federal funds rate less a survey-based measure of trend, over a sample of 1961-2007.

To evaluate power properties, DGPs 3 and 4 impose some parameter breaks on the specification of the bivariate DGP 1. In DGP 3, the breaks consist of one-time shifts in the intercept of the $y_{1,t}$ equation and the slope and intercept coefficients of the $y_{2,t}$ equation. The pre- and post-break coefficients are given in the last panel of Table 1; the error variance-covariance matrix is kept constant, at the setting used with DGP 1 (see above). The break

is imposed to occur at period $R + 1$, the date of the first out-of-sample forecast. In DGP 4, the single break takes the form of a shift in the error correlation ρ , from 0.5 to -0.5, at period $R + 1$. In this DGP, the other VAR parameters are stable over time, at the DGP 1 values given in Table 1.

In experiments with DGPs 1, 1G, 3, and 4, to keep the conditioning relatively simple and speed the Monte Carlo replications, we report results for forecast horizons of 1 and 2 periods ahead. In these cases, we forecast the first variable of the VAR in periods $t + 1$ and $t + 2$ (i.e., $y_{1,t+1}$ and $y_{1,t+2}$), conditional on the actual values of the second variable in those periods (i.e., $y_{2,t+1}$ and $y_{2,t+2}$). In experiments with DGP 2, we extend the conditioning horizon to 4 quarters ahead. In this case, we forecast variables $y_{1,t+\tau}$ and $y_{2,t+\tau}$ for $\tau = 1, \dots, 4$ conditional on $y_{3,t+\tau}$ equaling its actual values in periods $t + 1$ through $t + 4$ — that is, conditional on $\hat{y}_{3,t,\tau}^c = y_{3,t+\tau}$, $\tau = 1, \dots, 4$. For evaluating conditional forecasts produced under the policy shock approach, because 1-step ahead conditional forecasts obtained via the policy shock approach are exactly the same as the unconditional forecasts, we only report results for the multi-step horizons covered with each DGP.¹¹

4.2 Test implementation

For convenience, we list here the regressions and test statistics we use, referring to variable 1 of the VAR for convenience (in DGP 2, we consider forecasts of an additional variable):

$$\hat{\varepsilon}_{1,t,\tau}^i = \alpha_0 + e_{t,\tau}, \quad i = u, c, \quad \text{bias: } \alpha_0 \text{ } t\text{-stat} \quad (2)$$

$$\hat{\varepsilon}_{1,t,\tau}^i = \alpha_0 + \alpha_1 \hat{y}_{1,t,\tau}^i + e_{t,\tau}, \quad i = u, c, \quad \text{M-Z: } \alpha_1 \text{ } t\text{-stat} \quad (3)$$

$$\hat{\varepsilon}_{1,t,\tau}^c = \alpha_0 + \alpha_1 (\hat{y}_{1,t,\tau}^u - \hat{y}_{1,t,\tau}^c) + \alpha_2 \hat{y}_{1,t,\tau}^c + e_{t,\tau}, \quad \text{F-W: } \alpha_2 \text{ } t\text{-stat} \quad (4)$$

$$(\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 = \alpha_0 + e_{t,\tau}, \quad \text{MSE accuracy: } \alpha_0 = k(\hat{\phi}_T) \text{ } t\text{-stat} \quad (5)$$

To test bias, we regress (equation (2)) forecast errors (either unconditional or conditional) on a constant and form the t -statistic for the null of a coefficient of zero. To test efficiency, for all types of forecasts, we consider the Mincer-Zarnowitz efficiency test of equation (3). For conditional forecasts, we also consider the t -statistic of the coefficient on the conditional forecast in the Faust and Wright (2008) efficiency regression (4). Finally, we use the regression (5) to test whether the difference in MSEs for the unconditional and

¹¹In these DGPs, the equivalence of the 1-step conditional forecast obtained via the policy shock approach to the unconditional forecast arises because the “policy” variable is ordered last in the VAR.

conditional forecasts is different from the population level-value implied by the VAR’s parameters (where, as noted above, that population value is a function of the VAR’s slope coefficients and error variance-covariance matrix).

In all cases, the forecast error for horizon τ should follow an $\text{MA}(\tau - 1)$ process. At the 1-step horizon, we form test statistics using the simple OLS estimate of the variance. At longer horizons, we use a rectangular kernel, with $\tau - 1$ lags.¹²

4.3 Results

Tables 2 through 5 provide size results from Monte Carlo experiments with DGPs 1, 1G, and 2, in which the conditional forecasts are computed with the minimum-MSE approach.

These results indicate that comparing tests of bias, efficiency, and MSE accuracy against standard normal critical values will often be unreliable. More specifically:

- Under a recursive scheme, without correction for the effects of parameter estimation error, tests of bias — both unconditional and conditional — range from being about correctly sized (DGP 1) to modestly oversized (DGP 3). Oversizing is somewhat more likely at longer forecast horizons, probably because of difficulties with the finite-sample precision of HAC variance estimates.
- Under a recursive scheme, tests of efficiency range from being almost correctly sized (DGP 1, 1-step horizon) to significantly oversized (DGP 3). Size is comparable for the M-Z and F-W tests applied to conditional forecasts and the M-Z test applied to unconditional forecasts. The size distortions generally rise as the forecast horizon increases, likely reflecting imprecision in the HAC variance estimate. For example, for forecasts of $y_{2,t}$ in experiments with DGP 2 and $R, P = 100, 100$, the rejection rate for the normal-based M-Z efficiency test applied to conditional forecasts rises from 23.0 percent at the 1-step horizon to 29.0 percent at the 2-step horizon and 36.1 percent at the 4-step horizon.
- With recursively produced forecasts, the MSE accuracy test often, although not always, leads to significant undersizing. For example, with DGP 2, $R, P = 50, 100$, and the 1-step forecast horizon, the rejection rate for the MSE test is 3.5 percent for variable 1 and 14.3 percent for variable 2.

¹²In the rare occasions in which the result variance estimator is not positive semi-definite, we instead use the HAC estimator of Newey and West (1987), with $2(\tau - 1)$ lags.

- Under the rolling scheme, the failure to correct standard errors for the effects of parameter estimation error associated with forecast model estimation can have large consequences. In particular, rejection rates for bias tests are materially lower under the rolling scheme than the recursive, such that bias tests applied to rolling forecasts are often modestly or significantly undersized. In the experiments in which $P/R > 1$, rejection rates for efficiency tests are higher under the rolling scheme than the recursive, such that efficiency tests applied to rolling forecasts are often significantly oversized. Finally, tests of MSE accuracy in DGP 1 applied under the rolling scheme range from significantly to slightly undersized.
- Under the recursive scheme, conditional heteroskedasticity in the data-generating process does not materially affect the results described above. Broadly, size results under DGP 1G (Table 5) are fairly similar to those under DGP 1 (Table 2). The oversizing of efficiency tests is a little greater in the GARCH results than in the conditionally homoskedastic results, but otherwise results are fairly similar.

In comparison, conducting inference on the basis of our proposed bootstrap yields more reliable tests of bias, efficiency, and accuracy. More specifically:

- For both unconditional and conditional forecasts, the median rejection rate across all bias tests in Tables 2-5 is about 11.4 percent, with a minimum of about 9 percent and maximum of about 13-14 percent.
- For each type of efficiency test, the median rejection rate across all experiments in Tables 2-5 is a little less than 10 percent. In some settings, however, the tests can be either slightly-to-modestly undersized or slightly-to-modestly oversized. For example, applied to unconditional forecasts the M-Z test of efficiency has a minimum of 4.6 percent and maximum of 14.1 percent across experiments. Applied to conditional forecasts the M-Z test's rejection rate ranges from 5.9 percent to 14.3 percent.
- Size performance is more variable for the MSE test (unconditional versus minimum-MSE conditional) than the bias and efficiency tests. On average, the bootstrap is reasonably reliable: the median rejection rate across all MSE tests in Tables 2-5 is 8.9 percent. But the rejection rate ranges from 3.2 to 17.1 percent, depending on the experiment, variable, horizon, etc.

- With bootstrap inference, using a rolling scheme for estimation or a DGP with GARCH yields results similar to those obtained for recursive estimation of a DGP with conditionally homoskedastic innovations.

Tables 6 and 7 provide power results from Monte Carlo experiments with DGPs 3 and 4, in which the conditional forecasts are computed with the minimum-MSE approach and the forecasting scheme is recursive. We focus on power results obtained under bootstrap critical values, because using normal critical values (without correction for parameter estimation error) does not yield accurately sized tests. These experiments yield the following findings for power.

- With a break in VAR coefficients (DGP 3), tests for bias have relatively good power. For example, with $R, P = 100, 100$, the 2-step ahead rejection rate is 85.0 percent for the unconditional forecast and 74.8 percent for the conditional forecast. As this example indicates, the bias test for unconditional forecasts has somewhat more power than the bias test for conditional forecasts, likely because conditioning on the actual future path of $y_{2,t}$ reduces shifts in $y_{2,t}$ as a potential source of instability in the forecast of $y_{1,t}$.
- In the same DGP, tests for efficiency have at best modest power, with the M-Z test applied to unconditional forecasts having a little more power than the M-Z or F-W tests applied to conditional forecasts. In the same example ($R, P = 100, 100$ and a 2-step forecast horizon), the rejection rate is 17.3, 7.9, and 7.1 percent for the M-Z unconditional, M-Z conditional, and F-W conditional tests.
- Similarly, in the DGP with a break in VAR coefficients, the test of MSE accuracy has at least a little power. In the same example, the power of the MSE accuracy test is 14.9 percent (Table 6).
- In the DGP with a break in error correlations (Σ) instead of the VAR's slope coefficients, the power rankings of the tests are essentially reversed. In DGP 4, the MSE accuracy test has the best power, ranging from about 40 percent (with $R, P = 50, 150$) to as much as 94.5 percent ($R, P = 100, 50$, 1-step ahead forecast horizon). The M-Z efficiency test applied to conditional forecasts ranks second in power, yielding a rejection rate as high as 75.7 percent ($R, P = 100, 50$, 2-step ahead forecast horizon),

but also as low as 11.9 percent (with $R, P = 50, 150$, 1-step ahead forecast horizon). Neither the F-W test for the efficiency of conditional forecasts nor the M-Z test for the efficiency of unconditional forecasts have much power. Bias tests also lack power in the face of a shift in the error correlation (in Σ), likely due to the absence of any mean shifts that would lead to bias in forecasts.

Finally, Table 8 provides some size and power results for two-step ahead conditional forecasts produced under the policy shock approach.¹³ Broadly, these results are mostly, although not entirely, similar to the results for minimum-MSE conditional forecasts.

- In size experiments with DGP 1, the performance of the bias test for policy shock-based forecasts is about the same as for the unconditional and minimum-MSE conditional forecasts (compare Tables 2 and 8). Against standard normal critical values, the efficiency test rejection rates are modestly higher for policy shock-based forecasts than for the others. The oversizing of the M-Z efficiency test may not be surprising in light of the analytical results above that suggest that the M-Z efficiency test should reject with policy shock-conditioned forecasts. However, in these experiments, under bootstrap critical values, the oversizing goes away.
- In power experiments with DGP 3, the test for bias has more power than the tests for efficiency. With policy shock conditioning, the bias test has power comparable to the bias test applied to unconditional forecasts and above the power of the bias test applied to minimum-MSE conditional forecasts (Tables 2 and 8). Similarly, the efficiency tests have slightly to modestly more power with policy shock-conditioned forecasts than with the minimum-MSE forecasts.
- In power experiments with DGP 4, neither tests for bias nor tests for efficiency have much power. The primary difference with respect to the tests based on minimum-MSE conditional forecasts is that the M-Z efficiency test has little power, rather than modest to decent power, with policy shock-conditioned forecasts.

In summary, the Monte Carlo results are generally consistent with the analytical findings described above. With conditional forecasts, comparing tests of bias, efficiency, and MSE accuracy against standard normal critical values does not yield generally reliable results. A

¹³These results are based on exactly the same artificial data as the corresponding experiments in Tables 2, 6, and 7.

likely reason is that, in many cases, the asymptotic distributions of the tests are affected by parameter estimation error (from forecast model estimation). While the statistics of interest are normally distributed, parameter estimation error affects the appropriate variance in a way that is often very complicated. Our proposed (and easily implemented) bootstrap procedure yields tests with reasonably reliable size properties. As to power, our results show that shifts in error correlations will lead to rejections of MSE accuracy (unconditional versus conditional) and the efficiency of conditional forecasts, but not rejections of the efficiency of unconditional forecasts.

5 Empirical Application

In our empirical application, we examine forecasts of U.S. GDP growth, the unemployment rate, and core PCE inflation obtained with a BVAR of 22 variables. The model is patterned after a specification considered in Giannone, Lenza, and Primiceri (2012), using variables in levels or log levels. A number of studies (e.g., Banbura, Giannone, and Reichlin 2010 and Giannone, Lenza, and Primiceri 2012) have found that larger models often forecast more accurately than smaller models.

This section proceeds by detailing the model and our implementation (some additional details are included in Appendix 2), data and forecasting details, and results.

5.1 Model and implementation

The model we use is a BVAR(5) of the form:

$$Y_t = C + A(L)Y_{t-1} + \varepsilon_t,$$

in which all variables enter in levels or log levels. The table below lists the 22 variables included and their transformations.

The prior for the model takes the conjugate Normal-inverted Wishart form used in studies such as Banbura, Giannone, and Reichlin (2010), which yields a Normal-inverted Wishart posterior (Appendix 2 details the estimation steps). The priors on the VAR coefficients $A(L)$ include Minnesota-style unit root priors, sums of coefficient priors, and initial observations priors, patterned after Sims and Zha (1998), for example.

More specifically, in the Minnesota-style component of the prior, we impose the prior

Variables in large BVAR

variable	transformation	prior mean on AR(1) coefficient
real GDP	ln	1
real personal consumption expenditures (PCE)	ln	1
real business fixed investment (BFI)	ln	1
real residential investment (RESINV)	ln	1
industrial production (IP)	ln	1
capacity utilization in manufacturing (CU)	ln	1
total hours worked in the nonfarm business sector (HOURS)	ln	1
payroll employment (PAYROLLS)	ln	1
unemployment rate	level	1
consumer sentiment	level	1
spot commodity price index	ln	1
core PCE price index	ln	1
price index for gross private domestic investment	ln	1
GDP price index	ln	1
real hourly compensation in the nonfarm business sector	ln	1
federal funds rate	level	1
M2	ln	1
total reserves	ln	1
S&P 500 index of stock prices	ln	1
1-year (constant maturity) Treasury bond yield	level	1
5-year (constant maturity) Treasury bond yield	level	1
real effective exchange rate for major currencies	ln	1

expectation and standard deviation of the coefficient matrices to be:

$$E[A_k^{(ij)}] = \begin{cases} 1 & \text{if } i = j, k = 1 \\ 0 & \text{otherwise} \end{cases}, \text{ st. dev.}[A_k^{(ij)}] = \frac{\lambda_1 \sigma_i}{k \sigma_j}, k = 1, \dots, 5, \quad (6)$$

where $A_k^{(ij)}$ denotes the element in position (i, j) in the matrix A_k . For each variable's intercept contained in C we assume an informative prior with mean 0 and standard deviation $\lambda_0 \sigma_i$. The shrinkage parameter λ_1 measures the overall tightness of the prior: when $\lambda_1 \rightarrow 0$ the prior is imposed exactly and the data do not influence the estimates, while as $\lambda_1 \rightarrow \infty$ the prior becomes loose and the prior information does not influence the estimates, which will approach the standard *OLS* estimates. To set each scale parameter σ_i we follow common practice (see e.g. Litterman (1986) and Sims and Zha (1998)) and set it equal to the standard deviation of the residuals from a univariate autoregressive model.

Doan, et al. (1984) and Sims (1993) have proposed complementing standard Minnesota priors with additional priors which favor unit roots and cointegration, and introduce correlations in prior beliefs about the coefficients in a given equation. Accordingly, in our benchmark specification, we also include the ‘‘sum of coefficients’’ and ‘‘dummy initial observation’’ priors proposed in Doan et al. (1984) and Sims (1993), respectively. Both these priors can be implemented by augmenting the system with dummy observations. The details of our implementation of this priors (including the definition of hyperparameters λ_3 and λ_4) are contained in Appendix 2.

Reflecting common settings in the literature (e.g., Sims and Zha (1998)), we set the hyperparameters that govern the tightness of the prior as follows:

$$\lambda_0 = 1; \lambda_1 = 0.2; \lambda_3 = 1; \lambda_4 = 1.$$

Studies such as Carriero, Clark, and Marcellino (2012) and Giannone, Lenza, and Primiceri (2012) that have considered various methods for optimizing the tightness of the prior have found that, in forecast accuracy, the typical hyperparameter settings offer gains in forecast accuracy that are comparable to those obtained by optimized hyperparameter settings.

The prior specification is completed by choosing v_0 and S_0 , the prior degrees of freedom and scale matrix of the inverted Wishart prior for Σ , so that the prior expectation of Σ is equal to a fixed diagonal residual variance $E[\Sigma] = \text{diag}(\sigma_1^2, \dots, \sigma_n^2)$. In particular, following Kadiyala and Karlsson (1997), we set the diagonal elements of S_0 to $s_{0ii} = (v_0 - n - 1)\sigma_i^2$ and $v_0 = n + 2$.

To streamline computations, particularly with a bootstrap approach to inference, we compute forecasts from the BVAR without simulation. We use the posterior mean of coefficients and the error covariance matrix (Σ , which we use in forming conditional forecasts) to compute point forecasts based on just the posterior mean coefficients, iterating forward from 1-step ahead forecasts. Carriero, Clark, and Marcellino (2012) found point forecasts obtained with this approach to be essentially the same as point forecasts obtained from Monte Carlo simulation.

5.2 Data and sample

In generating and evaluating forecasts, we use data taken from the FAME database of the Federal Reserve Board. The quarterly data on industrial production, capacity utilization, payroll employment, the unemployment rate, consumer sentiment, commodity prices, interest rates, M2, reserves, stock prices, and the exchange rate are constructed as simple within-quarter averages of the source monthly data (in keeping with the practice of, e.g., Blue Chip and the Federal Reserve). All growth and inflation rates are measured as annualized log changes (from $t - 1$ to t).

In keeping with common practice in recent VAR forecasting analyses (e.g., Giannone, Lenza, and Primiceri 2012, Koop 2013) and with the FOMC’s forecast reporting practices, we provide results for a subset of the variables included in our models: GDP growth, the unemployment rate, and core PCE inflation. Unlike these studies, we do not report forecasts for the federal funds rate because we are interested in forecasts of the other variables conditional on particular paths of the federal funds rate.

We generate and evaluate forecasts for a sample of 1991 through 2007, at horizons of 1, 2, 4, and 8 quarters. We start forecasting in 1991 and not sooner because, through the 1980s, inflation was still trending down, presumably due to a deliberate disinflation effort by the Federal Reserve. We stop our forecast evaluation in 2007:Q4 to avoid possible complications of the zero lower bound constraints that became relevant in subsequent years.

In all cases, we estimate the model with a sample that starts in 1961:Q1. We begin by estimating with data from 1961:Q1 through 1990:Q4 and forming forecasts for 1991:Q1 through 1992:Q4. We then proceed by moving to a forecast origin of 1991:Q2, estimating the model with data from 1961:Q1 through 1991:Q1 and forming forecasts for 1991:Q2 through 1993:Q1. We proceed similarly through time, up through 2007:Q3, to obtain a sample of forecasts from 1991 through 2007. At each point in time, we produce conditional

forecasts based on the actual future path of the federal funds rate over the 8-step forecast horizon. In the presented results, in the interest of brevity, we focus on conditional forecasts produced with the minimum-MSE approach. Conditional forecasts based on policy shocks yield similar findings.

5.3 Results

Tables 9 and 10 provide the results of bias, efficiency, and MSE accuracy tests applied to growth, unemployment, and inflation forecasts. In particular, Table 9 reports test statistics (t -statistics) along with bootstrapped 10 percent critical values. Table 10 reports the MSEs of the unconditional and conditional forecasts, along with the t -test for MSE accuracy and its 10 percent critical values.

Overall, there is little evidence of bias in the growth, unemployment, or inflation forecasts, either unconditional or conditional. For GDP growth and unemployment, the forecast biases and test statistics provided in Table 9 are relatively small. Although the statistics are larger for inflation forecasts, they don't necessarily represent much evidence of bias. Whereas standard normal critical values would indicate the inflation forecasts — both unconditional and conditional — to be biased, the bootstrap critical values that our Monte Carlo analysis show to be more reliable indicate the null of no bias should not be rejected. Consider, for example, 4-quarter ahead unconditional forecasts of inflation. The t -statistic for the null of 0 bias in the unconditional forecast is -2.326, which would imply rejection of the null of no bias. However, the associated left tail bootstrap critical value (associated with a 2-sided significance level of 10%), is -3.198; based on the bootstrap critical values, the null of zero bias cannot be rejected. Therefore, over the 1991-2007 period, conditioning the forecasts on path of the federal funds rate that matches the actual path does not introduce any bias into the forecasts of inflation or the forecasts of GDP growth and unemployment.

There is considerably more evidence of inefficiency in the GDP growth and inflation forecasts — both unconditional and conditional — but not unemployment forecasts. At most horizons, all of the inefficiency tests reject the null of efficiency for growth and inflation forecasts. For example, in 4-step forecasts of GDP growth and inflation, the M-Z unconditional, M-Z conditional, and F-W conditional test statistics all fall outside the left tail of the bootstrap distribution.

Finally, the results in Table 10 indicate that the differences in MSE accuracy between the unconditional and minimum-MSE conditional forecasts are generally not significantly

different from what full sample estimates of the BVAR imply should obtain. (One other interesting finding in Table 10 is that, based on BVAR parameters, conditioning forecasts on the future path of interest rates should generally be expected to have modest effects on accuracy, except at the 8-step forecast horizon.) To be sure, there are differences in the actual MSEs that are somewhat different from the MSEs implied by the VAR’s parameters, but these differentials are not often statistically significant. Consider, for example, 4-quarter ahead forecasts of GDP growth. In the out-of-sample forecasts, the difference in MSEs (unconditional less minimum-MSE conditional) is 0.835. The BVAR, with full-sample parameter estimates, implies the difference in MSEs could be expected (in population) to be 0.322. However, the t -statistic for the equality of the sample difference and population-implied difference is just 1.085, well below the upper tail of the bootstrap distribution. The same outcome occurs with the other variables and forecast horizons, with the exception of a rejection of the null for inflation forecasts at the 8-step ahead horizon.

Overall, these results suggest some misspecification of the BVAR used to produce the forecasts of GDP growth, unemployment, and core inflation — both unconditional forecasts and forecasts conditioned on the actual path of the federal funds rate over a two year horizon. Although that misspecification doesn’t lead to significant bias in the forecasts or cause the difference in the accuracy of unconditional and conditional forecasts to be significantly different than the VAR’s parameters imply, it leads to a number of rejections of the efficiency of both unconditional and conditional forecasts.

6 Conclusions

Motivated by the common use of conditional forecasting in both practical forecasting and research, we develop and apply methods for the evaluation of conditional forecasts from VARs, using tests of bias, efficiency, and the MSE accuracy of conditional versus unconditional forecasts. More specifically, we provide analytical, Monte Carlo, and empirical evidence on tests of predictive ability for conditional forecasts from estimated models. Throughout, our intention is to provide usable metrics for evaluating conditional forecasts, in a general sense and in comparison to the accuracy of unconditional forecasts. To do so, we focus on particular forms of conditional forecasts for which interpretation of various null and alternative hypotheses is most straight-forward. In particular, in our analysis, we consider forecasts conditioned on actual future information on some variables in the VAR model.

For these tests, we establish asymptotic normality in the context of VAR-based conditional forecasts. Our results follow from an application of West (1996) and West and McCracken (1998) and as such establish the role of estimation error on the asymptotic distribution. As a practical matter, the standard errors can be quite complex and as such we suggest and consider a bootstrap approach to inference that is valid when even estimation error contributes to the asymptotic variance of the test statistic. Monte Carlo evidence suggests that the tests can be reasonably well sized in samples of the size often seen in macroeconomic applications.

Building on these results, in our application, we evaluate unconditional and conditional forecasts from a common macroeconomic Bayesian VAR. We produce unconditional and conditional forecasts of GDP growth, unemployment, and inflation, in which the conditional forecasts are based on the actual path of the short-term interest rate over an eight quarter forecast horizon. Using bootstrap critical values, we find evidence of model misspecification in the form of rejections of the efficiency of unconditional and conditional forecasts of GDP growth and inflation.

7 Appendix 1: Theory Details

In this section we provide proofs of the Theorems described in the text. In addition to the notation from Section 2, let $z_{t+r} = (x'_{t+r}, \dots, x'_t)'$ where $r = \max(\tau, m)$. Throughout we ignore the finite sample difference between P and $P - \tau + 1$.

Proof of Theorem 1: Given A1 - A5, the proof follows from Theorems 4.1 and 4.2 of West and McCracken (2008).

Proof of Theorem 2: Recall that $f_{t,\tau}(\hat{\phi}_t) = (\text{vech}_{-1}(\hat{g}_t \hat{g}'_t), \hat{g}'_t \hat{\varepsilon}_{1,t,\tau}^c)'$. Given Theorem 4.1 of West (1996), the result will follow if A1 - A5 are sufficient for Assumptions 1-4 (W1-W4) in West (1996) when applied to $f_{t,\tau}(\hat{\phi})$ and h_s . W2 and W4 follow immediately from A2 and A5. For W1 and W3 it's useful to recall that every element of $f_{t,\tau}(\hat{\phi}_t)$, $h_s(\phi)$, $\partial f_{t,\tau}(\hat{\phi}_t)/\partial \phi$, $\partial h_s(\phi)/\partial \phi$, $\partial^2 f_{t,\tau}(\hat{\phi}_t)/\partial \phi \partial \phi'$, and $\partial^2 h_s(\phi)/\partial \phi \partial \phi'$ are twice continuously differentiable functions of polynomials of ϕ and are quadratics of z_{t+r} . This follows from (i) the fact that the unconditional forecasts are iterated multistep forecasts from a VAR, (ii) the definition of minimum-MSE and policy shock-based conditioning, and (iii) the definition of $f_{t,\tau}(\hat{\phi}_t)$ and $h_s(\phi)$. As such, A1 and A3 suffice for W1. Finally, since (i) fourth order stationarity of w_t implies covariance stationarity of $f_{t,\tau}$, $\partial f_{t,\tau}/\partial \phi$ and h_s , (ii) mixing is preserved by finite dimensioned functions of mixing variables, and (iii) the existence of $8d$ moments for w_t implies the existence of $4d$ moments for $f_{t,\tau}$, $\partial f_{t,\tau}/\partial \phi$ and h_s we find that A1 and A4 suffice for W3 and the proof is complete.

Proof of Theorem 3: Define $f_{t,\tau}(\hat{\phi}_t)$ by rewriting the moment condition as $(\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k(\hat{\phi}_T) = f_{t,\tau}(\hat{\phi}_t) - (k(\hat{\phi}_T) - k)$. We provide the proof separately for the case when $\pi = 0$ and when $\pi > 0$.

(a) Let $\pi = 0$. Assumptions A2 and A4 imply $T^{1/2}(\hat{\phi}_T - \phi) \rightarrow^d N(0, BS_{hh}B')$. Since $k(\phi)$ is continuously differentiable in ϕ , the Delta method implies $T^{1/2}(k(\hat{\phi}_T) - k) \rightarrow^d N(0, KBS_{hh}B'K')$. Since $\pi = 0$ implies $T/P = o(1)$, $P^{1/2}(k(\hat{\phi}_T) - k) = o_p(1)$ and hence $P^{-1/2} \sum_{t=R}^{T-\tau} ((\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k(\hat{\phi}_T)) = P^{-1/2} \sum_{t=R}^{T-\tau} ((\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k) + o_p(1)$. The result will follow if A1 - A5 are sufficient for assumptions W1-W4 in West (1996) when applied to $f_{t,\tau}(\hat{\phi})$ and h_s . Since the arguments used in the proof of Theorem 2 are applicable when $f_{t,\tau}(\hat{\phi}_t) = (\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k$, the result follows from Theorem 4.1 of West (1996).

(b) Let $\pi > 0$. From the proof of (a) it's clear that $P^{1/2}(k(\hat{\phi}_T) - k) = KB(P^{1/2}H(T)) + o_p(1) \rightarrow^d N(0, \frac{\pi}{1+\pi}KBS_{hh}B'K')$. In addition, Lemma 4.1 and Theorem 4.1 of West (1996)

imply

$$\begin{aligned} P^{-1/2} \sum_{t=R}^{T-\tau} ((\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k) &= P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau} + FB(P^{-1/2} \sum_{t=R}^{T-\tau} H(t)) + o_p(1) \\ &\rightarrow {}^dN(0, S_{ff} + 2\lambda_{fh}FB S'_{fh} + \lambda_{hh}FB S_{hh}B'F'). \end{aligned}$$

Together these imply

$$\begin{aligned} P^{-1/2} \sum_{t=R}^{T-\tau} ((\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k(\hat{\phi}_T)) &= P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau} + FB(P^{-1/2} \sum_{t=R}^{T-\tau} H(t)) \\ &\quad - KB(P^{1/2}H(T)) + o_p(1) \rightarrow {}^dN(0, \Omega_0) \end{aligned}$$

for some asymptotic variance Ω_0 . The result will follow if we can establish that $\Omega_0 = \Omega$ as described in the text. To do this note that we can rewrite $P^{-1/2} \sum_{t=R}^{T-\tau} ((\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k(\hat{\phi}_T))$ as

$$P^{-1/2} \sum_{t=R}^{T-\tau} ((\hat{\varepsilon}_{1,t,\tau}^c)^2 - (\hat{\varepsilon}_{1,t,\tau}^u)^2 - k(\hat{\phi}_T)) = [I, FB, -KB] \begin{pmatrix} P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau} \\ P^{-1/2} \sum_{t=R}^{T-\tau} H(t) \\ P^{1/2}H(T) \end{pmatrix} + o_p(1)$$

and hence

$$\Omega_0 = [I, FB, -KB] \lim_{P,R \rightarrow \infty} \text{Var}(P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}, P^{-1/2} \sum_{t=R}^{T-\tau} H(t)', P^{1/2}H(T)') [I, FB, -KB]'.$$

Straightforward algebra reveals that $\Omega_0 = \Omega$ if (i) $\lim_{P,R \rightarrow \infty} \text{Cov}(P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}, P^{1/2}H(T)') = \frac{\pi}{1+\pi} S'_{fh}$ and (ii) $\lim_{P,R \rightarrow \infty} \text{Cov}(P^{-1/2} \sum_{t=R}^{T-\tau} H(t), P^{1/2}H(T)') = \frac{\pi}{1+\pi} S_{hh}$. We establish both of these below accounting for whether the rolling or recursive schemes are used.

(b-i) Note that $P^{1/2}H(T) = \frac{(PR)^{1/2}}{T} (R^{-1/2} \sum_{s=1}^{R-1} h_{s+1}) + \frac{P}{T} (P^{-1/2} \sum_{s=R}^{T-1} h_{s+1})$. Given A4, it is straightforward to show that $\lim_{P,R \rightarrow \infty} \text{Cov}(P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}, R^{-1/2} \sum_{s=1}^{R-1} h'_{s+1}) = 0$. In addition, since A5 implies $\frac{(PR)^{1/2}}{T} = O(1)$, we immediately find that

$$\begin{aligned} \lim_{P,R \rightarrow \infty} \text{Cov}(P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}, P^{1/2}H(T)') &= \lim_{P,R \rightarrow \infty} \frac{P}{T} \text{Cov}(P^{-1/2} \sum_{t=R}^{T-\tau} f_{t,\tau}, P^{-1/2} \sum_{s=R}^{T-1} h'_{s+1}) \\ &= \frac{\pi}{1+\pi} S'_{fh} \end{aligned}$$

and we obtain the desired result. Note that this does not depend on whether the recursive or rolling scheme is used.

(b-ii) We do the decomposition three distinct ways: once for the recursive, once for the rolling with $P < R$, and once for the rolling with $P \geq R$.

(b-ii-recursive) Let $a_{R,s} = \sum_{j=0}^{P-1} (R+s+j)^{-1}$ and note that $P^{-1/2} \sum_{t=R}^{T-1} H(t)$ equals $(R/P)^{1/2} a_{R,0} (R^{-1/2} \sum_{s=1}^{R-1} h_{s+1}) + P^{-1/2} \sum_{s=R+1}^{T-1} a_{R,s} h_{s+1}$. And as above, $P^{1/2} H(T) = \frac{(PR)^{1/2}}{T} (R^{-1/2} \sum_{s=1}^{R-1} h_{s+1}) + \frac{P}{T} (P^{-1/2} \sum_{s=R}^{T-1} h_{s+1})$. Given A4, showing that $\lim_{P,R \rightarrow \infty} Cov(R^{-1/2} \sum_{s=1}^{R-1} h_{s+1}, P^{-1/2} \sum_{s=R}^{T-1} h_{s+1})$ and $\lim_{P,R \rightarrow \infty} Cov(R^{-1/2} \sum_{s=1}^{R-1} h_{s+1}, P^{-1/2} \sum_{s=R}^{T-1} a_{R,s} h_{s+1})$ are both zero is straightforward. In addition, since A5 implies $a_{R,0} \sim \ln(1+\pi)$ and $\frac{(PR)^{1/2}}{T}$ are $O(1)$ we find that

$$\begin{aligned} \lim_{P,R \rightarrow \infty} Cov(P^{-1/2} \sum_{t=R}^{T-\tau} H(t), P^{1/2} H(T)) &= \lim_{P,R \rightarrow \infty} \frac{R}{T} a_{R,0} Cov(R^{-1/2} \sum_{s=1}^{R-1} h_{s+1}, R^{-1/2} \sum_{s=1}^{R-1} h'_{s+1}) \\ &+ \lim_{P,R \rightarrow \infty} \frac{P}{T} Cov(P^{-1/2} \sum_{s=R}^{T-1} a_{R,s} h_{s+1}, P^{-1/2} \sum_{s=R}^{T-1} h'_{s+1}). \end{aligned}$$

Since $\lim_{P,R \rightarrow \infty} Cov(R^{-1/2} \sum_{s=1}^{R-1} h_{s+1}, R^{-1/2} \sum_{s=1}^{R-1} h_{s+1}) = \lim_{P,R \rightarrow \infty} Var(R^{-1/2} \sum_{s=1}^{R-1} h_{s+1})$ we immediately find that the first right-hand side term equals $(1+\pi)^{-1} \ln(1+\pi)$. That the second right-hand side term equals $\frac{\pi}{1+\pi} (1 - \pi^{-1} \ln(1+\pi))$ is delineated in the proof of Lemma A6 in West (1996). Adding the two pieces together provides the desired result.

(b-ii-rolling, $P < R$) Define $\sum_{t=R}^{T-1} H(t) = R^{-1} \sum_{s=1}^{P-1} s h_s + \frac{P}{R} \sum_{s=P}^R h_s + R^{-1} \sum_{s=R+1}^{T-1} (P-s-R) h_s = A_1 + A_2 + A_3$ and $H(T) = T^{-1} \sum_{s=1}^{P-1} h_s + T^{-1} \sum_{s=P}^R h_s + T^{-1} \sum_{s=R+1}^{T-1} h_s = B_1 + B_2 + B_3$. Given A4, it is straightforward to show $\lim_{P,R \rightarrow \infty} Cov(A_i, B'_j) = 0$ all $i \neq j$ and hence $\lim_{P,R \rightarrow \infty} Cov(P^{-1/2} \sum_{t=R}^{T-\tau} H(t), P^{1/2} H(T)') = \sum_{j=1}^3 \lim_{P,R \rightarrow \infty} Cov(A_j, B'_j)$. For the second term¹⁴ it's clear that $\lim_{P,R \rightarrow \infty} Cov(A_2, B'_2) = \lim_{P,R \rightarrow \infty} \frac{P(R-P)}{RT} Cov((R-P)^{-1/2} \sum_{s=P}^R h_s, (R-P)^{-1/2} \sum_{s=P}^R h'_s) = \frac{\pi}{1+\pi} (1-\pi) S_{hh}$.

For the first and third terms a bit more detail is needed. For ease of presentation let h_s be a scalar and define $\gamma_j = E h_s h_{s-j}$, $d_j = \sum_{i=1}^{P-1-j} i$, and $c_j = \sum_{i=1}^{P-1-j} (P-i)$. Direct multiplication and taking expectations implies $\lim_{P,R \rightarrow \infty} Cov(A_1, B_1) = (RT)^{-1} [\sum_{j=1}^{P-2} \gamma_j d_j + \sum_{j=1}^{P-2} \gamma_j c_j + \gamma_0 d_0]$. A5 and straightforward algebra imply $(RT)^{-1} d_0 = \frac{P^2}{RT} (P^{-2} \sum_{i=1}^{P-2} i) \rightarrow \frac{\pi^2}{2(1+\pi)}$. Since $\gamma_0 + 2 \sum_{j=1}^{P-2} \gamma_j \rightarrow S_{hh}$ the result will follow if $P^{-2} |\sum_{j=1}^{P-2} \gamma_j (d_j - d_0)|$ and $P^{-2} |\sum_{j=1}^{P-2} \gamma_j (c_j - d_0)|$ are both $o(1)$. We show the former, the proof of the latter is very similar. To do so note that $d_j \geq \int_1^{P-1} (x-j) dx$ and $d_0 \leq \int_0^P x dx$. Integrating we obtain $|d_j - d_0| \leq |P + j(P - \frac{3}{2})|$ and hence $P^{-2} |\sum_{j=1}^{P-2} \gamma_j (d_j - d_0)| \leq P^{-2} \sum_{j=1}^{P-2} |\gamma_j| |P + j(P - \frac{3}{2})| \leq P^{-1} \sum_{j=1}^{P-2} |\gamma_j| + P^{-1} \sum_{j=1}^{P-2} |\gamma_j| |j|$. Since A4 implies $\sum_{j=1}^{P-2} |\gamma_j|$ and $\sum_{j=1}^{P-2} |\gamma_j| |j|$ are $O(1)$ it is clear that $P^{-2} |\sum_{j=1}^{P-2} \gamma_j (d_j - d_0)| = o(1)$ and we

¹⁴Here we treat the case in which $|P-R|$ diverges as $P, R \rightarrow \infty$. When the difference is finite the second term is trivially $o_p(1)$. The distinction does not effect the ultimate formula for the asymptotic variance since, if $|P-R|$ diverges and $\pi = 1$, we still obtain the result that the second term is $o_p(1)$ (because the asymptotic variance of this term is zero).

conclude that $\lim_{P,R \rightarrow \infty} Cov(A_1, B_1) = \frac{\pi^2}{2(1+\pi)} S_{hh}$.

Moving to the third term, direct multiplication and taking expectations implies $\lim_{P,R \rightarrow \infty} Cov(A_3, B_3) = (RT)^{-1} [\sum_{j=1}^{P-2} \gamma_j d_j + \sum_{j=1}^{P-2} \gamma_j c_j + \gamma_0 d_0]$ if we redefine $c_j = \sum_{i=2}^{P-j} i$. Unsurprisingly, since this expansion is nearly identical to that for the first term, nearly identical arguments imply $\lim_{P,R \rightarrow \infty} Cov(A_3, B_3) = \frac{\pi^2}{2(1+\pi)} S_{hh}$.

Finally, if we add the three terms together we find that $\sum_{j=1}^3 \lim_{P,R \rightarrow \infty} Cov(A_j, B'_j) = \frac{\pi^2}{2(1+\pi)} S_{hh} + \frac{\pi}{1+\pi} (1-\pi) S_{hh} + \frac{\pi^2}{2(1+\pi)} S_{hh} = \frac{\pi}{1+\pi} S_{hh}$ and the proof is complete.

(b-ii-rolling, $P \geq R$) Define $\sum_{t=R}^{T-1} H(t) = R^{-1} \sum_{s=1}^R sh_s + \sum_{s=R+1}^P h_s + R^{-1} \sum_{s=1}^{R-1} sh_{T-s} = A_1 + A_2 + A_3$ and $H(T) = T^{-1} \sum_{s=1}^R h_s + T^{-1} \sum_{s=R+1}^P h_s + T^{-1} \sum_{s=1}^{R-1} h_{T-s} = B_1 + B_2 + B_3$. Given A4, it is straightforward to show $\lim_{P,R \rightarrow \infty} Cov(A_i, B'_j) = 0$ all $i \neq j$ and hence $\lim_{P,R \rightarrow \infty} Cov(P^{-1/2} \sum_{t=R}^{T-\tau} H(t), P^{1/2} H(T)') = \sum_{j=1}^3 \lim_{P,R \rightarrow \infty} Cov(A_j, B'_j)$. For the second term¹⁵ it's clear that $\lim_{P,R \rightarrow \infty} Cov(A_2, B'_2) = \lim_{P,R \rightarrow \infty} \frac{(P-R)}{T} Cov((P-R)^{-1/2} \sum_{s=R+1}^P h_s, (P-R)^{-1/2} \sum_{s=R+1}^P h'_s) = \frac{\pi-1}{1+\pi} S_{hh}$.

Once again, for the first and third terms a bit more detail is needed. For ease of presentation let h_s be a scalar and define $\gamma_j = Eh_s h_{s-j}$, $d_j = \sum_{i=1}^{R-j} i$, and $c_j = \sum_{i=j+1}^R i$. Direct multiplication and taking expectations implies $\lim_{P,R \rightarrow \infty} Cov(A_1, B_1) = (RT)^{-1} [\sum_{j=1}^{R-1} \gamma_j d_j + \sum_{j=1}^{R-1} \gamma_j c_j + \gamma_0 d_0]$. A5 and straightforward algebra imply $(RT)^{-1} d_0 = \frac{R^2}{RT} (R^{-2} \sum_{i=1}^{R-1} i) \rightarrow \frac{1}{2(1+\pi)}$. Since $\gamma_0 + 2 \sum_{j=1}^{R-1} \gamma_j \rightarrow S_{hh}$ the result will follow if $R^{-2} |\sum_{j=1}^{R-1} \gamma_j (d_j - d_0)|$ and $R^{-2} |\sum_{j=1}^{R-1} \gamma_j (c_j - d_0)|$ are both $o(1)$. We show the former, the proof of the latter is very similar. To do so note that $d_j \geq \int_1^{R-1} (x-j) dx$ and $d_0 \leq \int_0^R x dx$. Integrating we obtain $|d_j - d_0| \leq |R + j(R-2)|$ and hence $R^{-2} |\sum_{j=1}^{R-1} \gamma_j (d_j - d_0)| \leq R^{-2} \sum_{j=1}^{R-1} |\gamma_j| |R + j(R-2)| \leq R^{-1} \sum_{j=1}^{R-1} |\gamma_j| + R^{-1} \sum_{j=1}^{R-1} |\gamma_j| |j|$. Since A4 implies $\sum_{j=1}^{R-1} |\gamma_j|$ and $\sum_{j=1}^{R-1} |\gamma_j| |j|$ are $O(1)$ it is clear that $R^{-2} |\sum_{j=1}^{R-1} \gamma_j (d_j - d_0)| = o(1)$ and we conclude that $\lim_{P,R \rightarrow \infty} Cov(A_1, B_1) = \frac{1}{2(1+\pi)} S_{hh}$.

Moving to the third term, we again find that $\lim_{P,R \rightarrow \infty} Cov(A_3, B_3) = (RT)^{-1} [\sum_{j=1}^{R-2} \gamma_j d_j + \sum_{j=1}^{R-2} \gamma_j c_j + \gamma_0 d_0]$ if we redefine $d_j = \sum_{i=1}^{R-1-j} i$ and $c_j = \sum_{i=j+1}^{R-1} i$. Unsurprisingly, since this expansion is nearly identical to that for the first term, nearly identical arguments imply $\lim_{P,R \rightarrow \infty} Cov(A_3, B_3) = \frac{1}{2(1+\pi)} S_{hh}$.

Finally, if we add the three terms together we find that $\sum_{j=1}^3 \lim_{P,R \rightarrow \infty} Cov(A_j, B'_j) = \frac{1}{2(1+\pi)} S_{hh} + \frac{\pi-1}{1+\pi} S_{hh} + \frac{1}{2(1+\pi)} S_{hh} = \frac{\pi}{1+\pi} S_{hh}$ and the proof is complete.

¹⁵See the previous footnote.

8 Appendix 2: BVAR Details

This appendix provides additional detail on the procedure for estimating the BVAR used in the application and on the prior used in estimation.

8.1 Estimation procedure for BVAR

By grouping the coefficient matrices in the $n \times M$ matrix $A' = [C \ A_1 \ \dots \ A_l]$ and defining $x_t = (1 \ y'_{t-1} \ \dots \ y'_{t-l})'$ as a vector containing an intercept and l lags of y_t , the VAR can be written as:

$$y_t = \Lambda' x_t + \varepsilon_t. \quad (7)$$

An even more compact notation is:

$$Y = X\Lambda + E, \quad (8)$$

where $Y = [y_1, \dots, y_T]'$, $X = [x_1, \dots, x_T]'$, and $E = [\varepsilon_1, \dots, \varepsilon_T]'$ are, respectively, $T \times n$, $T \times M$ and $T \times n$ matrices.

We use the conjugate Normal-inverted Wishart prior:

$$\Lambda | \Sigma \sim N(\Lambda_0, \Sigma \otimes \Omega_0), \quad \Sigma \sim IW(S_0, v_0). \quad (9)$$

As the N-IW prior is conjugate, the conditional posterior distribution of this model is also N-IW (Zellner 1971):

$$\Lambda | \Sigma, Y \sim N(\bar{\Lambda}, \Sigma \otimes \bar{\Omega}), \quad \Sigma | Y \sim IW(\bar{S}, \bar{v}). \quad (10)$$

Defining $\hat{\Lambda}$ and \hat{E} as the OLS estimates, we have that $\bar{\Lambda} = (\Omega_0^{-1} + X'X)^{-1}(\Omega_0^{-1}\Lambda_0 + X'Y)$, $\bar{\Omega} = (\Omega_0^{-1} + X'X)^{-1}$, $\bar{v} = v_0 + T$, and $\bar{S} = \Lambda_0 + \hat{E}'\hat{E} + \hat{\Lambda}'X'X\hat{\Lambda} + \Lambda_0'\Omega_0^{-1}\Lambda_0 - \bar{\Lambda}'\bar{\Omega}^{-1}\bar{\Lambda}$. Sources such as Kadiyala and Karlsson (1997) and Banbura, Giannone, and Reichlin (2010) provide additional detail on the N-IW prior and posterior.

8.2 Prior for the BVAR

The ‘‘sum of coefficients’’ prior expresses a belief that when the average of lagged values of a variable is at some level \bar{y}_{0i} , that same value \bar{y}_{0i} is likely to be a good forecast of future observations, and it is implemented by augmenting the system in (8) with the dummy observations Y_{d_1} and X_{d_1} with generic elements:

$$y_d(i, j) = \begin{cases} \bar{y}_{0i}/\lambda_3 & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}; \quad x_d(i, s) = \begin{cases} \bar{y}_{0i}/\lambda_3 & \text{if } i = j, s < M \\ 0 & \text{otherwise,} \end{cases} \quad (11)$$

where i and j go from 1 to n while s goes from 1 to M . When $\lambda_3 \rightarrow 0$ the model tends to a form that can be expressed entirely in terms of differenced data, in which case there are as many unit roots as variables and there is no cointegration.

The “dummy initial observation” prior introduces a single dummy observation such that all values of all variables are set equal to the corresponding averages of initial conditions up to a scaling factor ($1/\lambda_4$). It is implemented by adding to the system in (8) the dummy variables Y_{d_2} and X_{d_2} with generic elements:

$$y_d(j) = \bar{y}_{0j}/\lambda_4; \quad x_d(s) = \begin{cases} \bar{y}_{0j}/\lambda_4 & \text{for } s < M \\ 1/\lambda_4 & \text{for } s = M, \end{cases} \quad (12)$$

where j goes from 1 to n while s goes from 1 to M . As $\lambda_4 \rightarrow 0$ the model tends to a form in which either all variables are stationary with means equal to the sample averages of the initial conditions, or there are unit root components without drift terms, which is consistent with cointegration.

References

- Aastveit, Knut Are, Andrea Carriero, Todd E. Clark, and Massimiliano Marcellino (2014), "Have Standard VARs Remained Stable Since the Crisis?" Federal Reserve Bank of Cleveland Working Paper Working Paper 14-11.
- Andrews, Donald W.K., and J. Christopher Monahan (1992), "An Improved Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimator," *Econometrica* 60, 953-966.
- Banbura, Michele, Domenico Giannone, and Lucrezia Reichlin (2010), "Large Bayesian Vector Autoregressions," *Journal of Applied Econometrics* 25, 71-92.
- Banbura, M., Giannone, D., and Reichlin, L., 2014. Conditional Forecasts and Scenario Analysis with Vector Autoregressions for Large Cross-Sections, *International Journal of Forecasting*, forthcoming.
- Baumeister, Christiane, and Lutz Kilian (2013), "Real-Time Analysis of Oil Price Risks Using Forecast Scenarios," *IMF Economic Review* 62, April 2014, 119-145.
- Bernanke, Ben S., and Alan S. Blinder (1992), "The Federal Funds Rate and the Channels of Monetary Transmission," *American Economic Review* 82, 901-921.
- Bruggemann, Ralf, Carsten Jentsch, and Carsten Trenkler (2014), "Inference in VARs with Conditional Heteroskedasticity of Unknown Form," manuscript, University of Mannheim.
- Carriero Andrea, Todd E. Clark, and Massimiliano Marcellino (2012), "Bayesian VARs: Specification Choices and Forecast Accuracy," *Journal of Applied Econometrics*, forthcoming.
- Christiano, Lawrence J., Martin Eichenbaum, and Charles Evans (1996), "The Effects of Monetary Policy Shocks: Evidence from the Flow of Funds," *Review of Economics and Statistics* 78, 16-34.
- Clarida, Richard H., and Coyle, Diane (1984), "Conditional Projection by Means of Kalman Filtering." Mimeo, National Bureau of Economic Research.
- Clark, Todd E., and Taeyoung Doh (2014), "Evaluating Alternative Models of Trend Inflation," *International Journal of Forecasting* 30, 426-448.
- Clark, Todd E., and Michael W. McCracken (2001), "Tests of Equal Forecast Accuracy and Encompassing for Nested Models," *Journal of Econometrics* 105, 85-110.
- Clark, Todd E., and Michael W. McCracken (2005), "Evaluating Direct Multistep Forecasts," *Econometric Reviews* 24, 369-404.
- Clark, Todd E., and Michael W. McCracken (2013), "Evaluating the Accuracy of Forecasts from Vector Autoregressions," in *Advances in Econometrics: VAR models in Macroeconomics — New Developments and Applications*, Thomas Fomby, Lutz Kilian, and Anthony Murphy (eds.), Emerald, 117-168.
- Del Negro, Marco, and Frank Schorfheide (2013), "DSGE Model-Based Forecasting," in G. Elliott and A. Timmermann (eds.), *Handbook of Economic Forecasting*, vol. 2, Amster-

dam: North-Holland.

- Diebold, Francis X., and Roberto S. Mariano (1995), "Comparing Predictive Accuracy," *Journal of Business and Economic Statistics* 13, 253-263.
- Doan, T., Litterman, R., and Sims, C. (1984), "Forecasting and Conditional Projection Using Realistic Prior Distributions," *Econometric Reviews* 3, 1-100.
- Faust, Jon, and Jonathan H. Wright (2008), "Efficient Forecast Tests for Conditional Policy Forecasts," *Journal of Econometrics* 146, 293-303.
- Faust, Jon, and Jonathan H. Wright (2013), "Forecasting Inflation," in *Handbook of Economic Forecasting*, volume 2 (North Holland).
- Gavin, William T. (2003), "FOMC Forecast: Is All the Information in the Central Tendency?" *Federal Reserve Bank of St. Louis Review*, May/June, 85, 27-48.
- Gavin, William T. and Mandal, Rachel J. (2003), "Evaluating FOMC Forecasts," *International Journal of Forecasting* 19, 655-67.
- Gavin, William T. and Geetanjali Pande (2008), "FOMC Consensus Forecasts," *Federal Reserve Bank of St. Louis Review*, May/June, 90 (Part 1), 146-163.
- Giacomini, Rafaella, and Halbert White (2006), "Tests of Conditional Predictive Ability," *Econometrica* 74, 1545-1578.
- Giannone, Domenico, Michele Lenza, and Giorgio Primiceri (2012), "Prior Selection for Vector Autoregressions," *Review of Economics and Statistics*, forthcoming.
- Giannone, D., Lenza, M., Momferatou, D., and Onorante L. (2014), "Short-Term Inflation Projections: A Bayesian Vector Autoregressive Approach," *International Journal of Forecasting* 30, 635-644.
- Goncalves, Silvia, and Lutz Kilian (2004), "Bootstrapping Autoregressions with Conditional Heteroskedasticity of Unknown Form," *Journal of Econometrics* 123, 89-120.
- Herbst, Edward, and Frank Schorfheide (2012), "Evaluating DSGE Model Forecasts of Comovements," *Journal of Econometrics* 171, 152-166.
- Jarocinski, Marek (2010), "Conditional Forecasts and Uncertainty about Forecast Revisions in Vector Autoregressions," *Economics Letters* 108, 257-259.
- Joutz, Fred and Herman Stekler (2000), "An Evaluation of the Predictions of the Federal Reserve," *International Journal of Forecasting* 16, 17-38.
- Kadiyala, K. Rao, and Sune Karlsson (1997), "Numerical Methods for Estimation and Inference in Bayesian VAR-Models," *Journal of Applied Econometrics* 12, 99-132.
- Kilian, Lutz (1999), "Exchange Rates and Monetary Fundamentals: What Do We Learn from Long-Horizon Regressions?" *Journal of Applied Econometrics* 14, 491-510.
- Koop, Gary (2013), "Forecasting with Medium and Large Bayesian VARs," *Journal of Applied Econometrics* 28, 177-203.
- Litterman, Robert B. (1986), "Forecasting with Bayesian Vector Autoregressions-Five Years

- of Experience," *Journal of Business and Economic Statistics* 4, 25-38.
- McCracken, Michael W. (2004), "Parameter Estimation Error and Tests of Equal Forecast Accuracy Between Non-nested Models," *International Journal of Forecasting* 20, 503-514.
- Meyer, Brent, and Saeed Zaman (2013), "It's Not Just for Inflation: The Usefulness of the Median CPI in BVAR Forecasting," Federal Reserve Bank of Cleveland Working Paper Working Paper 13-03.
- Mincer, J. and Victor Zarnowitz (1969), "The Evaluation of Economic Forecasts," in J. Mincer (ed.), *Economic Forecasts and Expectations*, 3-46, National Bureau of Economic Research.
- Newey, Whitney K., and Daniel L. McFadden (1994), "Large Sample Estimation and Hypothesis Testing," in R.F. Engle and D.L. Newey (eds.) *Handbook of Econometrics*, 4, 2113-2247, Elsevier Science.
- Newey, Whitney K., and Kenneth D. West (1987), "A Simple, Positive Semi-definite, Heteroskedasticity and Autocorrelation Consistent Covariance Matrix," *Econometrica* 55, 703-708.
- Patton, Andrew J., and Allan Timmermann (2012), "Forecast Rationality Tests Based on Multi-Horizon Bounds," *Journal of Business and Economic Statistics* 30, 1-17.
- Reifschneider, David, and Peter Tulip (2007), "Gauging the Uncertainty of the Economic Outlook from Historical Forecasting Errors," *FEDS WP* 2007-60.
- Romer, Christina D., and David H. Romer (2008), "The FOMC versus the Staff: Where Can Monetary Policymakers Add Value?" *American Economic Review* 98, 230-235.
- Schorfheide, Frank, and Dongho Song (2013), "Real Time Forecasting with a Mixed-Frequency VAR," NBER Working Paper 19712.
- Sims, Christopher A. (1982), "Policy Analysis with Econometric Models," *Brookings Papers on Economic Activity*, No. 1, 107-152.
- Sims, C., and Zha, T. (1998), "Bayesian Methods for Dynamic Multivariate Models," *International Economic Review* 39, 949-968.
- Stine, Robert A. (1987), "Estimating Properties of Autoregressive Forecasts," *Journal of the American Statistical Association* 82, 1072-1078.
- Waggoner D.F., and Zha T. (1999), "Conditional Forecasts in Dynamic Multivariate Models," *The Review of Economics and Statistics* 81, 639-651
- West, Kenneth D. (1996), "Asymptotic Inference About Predictive Ability," *Econometrica* 64, 1067-1084.
- West, Kenneth D., and Michael W. McCracken (1998), "Regression-based Tests of Predictive Ability," *International Economic Review* 39, 817-840.
- Zellner, Arnold (1971), *An Introduction to Bayesian Inference in Econometrics*, J. Wiley and Sons, Inc., New York.

Table 1. Monte Carlo DGP coefficients

explanatory variable	$y_{1,t}$ equation	$y_{2,t}$ equation	$y_{3,t}$ equation
DGP 1 (size)			
$y_{1,t-1}$	0.50	0.00	
$y_{2,t-1}$	0.10	0.80	
intercept	0.00	0.00	
DGP 2 (size)			
$y_{1,t-1}$	0.234	0.029	0.059
$y_{1,t-2}$	0.164	-0.039	0.031
$y_{2,t-1}$	-0.134	0.575	0.038
$y_{2,t-2}$	-0.150	0.138	0.019
$y_{3,t-1}$	-0.057	0.200	1.006
$y_{3,t-2}$	-0.165	-0.184	-0.087
intercept	2.425	0.054	-0.110
DGP 3 (power)			
$y_{1,t-1}$, pre-break	0.50	0.00	
$y_{1,t-1}$, post-break	0.50	0.25	
$y_{2,t-1}$, pre-break	0.10	0.80	
$y_{2,t-1}$, post-break	0.10	0.40	
intercept, pre-break	0.00	0.00	
intercept, post-break	0.50	0.40	

Notes:

1. The table provides the coefficients of Monte Carlo DGPs 1-3. Other details of DGPs 1G and 4 are given in section 4.1.
2. The variance-covariance matrix of innovations and other aspects of the Monte Carlo design are described in section 4.1.

Table 2: Monte Carlo Results on Size, Minimum-MSE Conditioning, DGP 1
(nominal size = 10%)

1-step horizon					
<i>test</i>	<i>source of critical values</i>	<i>R=50</i> <i>P=100</i>	<i>R=50</i> <i>P=150</i>	<i>R=100</i> <i>P=50</i>	<i>R=100</i> <i>P=100</i>
bias, uncond.	Normal	0.115	0.132	0.118	0.123
bias, condit.	Normal	0.119	0.121	0.119	0.116
M-Z efficiency, uncond.	Normal	0.142	0.117	0.141	0.113
M-Z efficiency, condit.	Normal	0.170	0.132	0.146	0.114
F-W efficiency, condit.	Normal	0.147	0.102	0.126	0.101
equal MSE	Normal	0.008	0.005	0.057	0.026
bias, uncond.	bootstrap	0.118	0.122	0.128	0.114
bias, condit.	bootstrap	0.116	0.114	0.129	0.105
M-Z efficiency, uncond.	bootstrap	0.128	0.103	0.130	0.104
M-Z efficiency, condit.	bootstrap	0.121	0.095	0.132	0.103
F-W efficiency, condit.	bootstrap	0.129	0.104	0.127	0.100
equal MSE	bootstrap	0.107	0.100	0.135	0.098
2-step horizon					
bias, uncond.	Normal	0.116	0.126	0.130	0.121
bias, condit.	Normal	0.118	0.117	0.131	0.117
M-Z efficiency, uncond.	Normal	0.257	0.227	0.264	0.182
M-Z efficiency, condit.	Normal	0.205	0.174	0.196	0.148
F-W efficiency, condit.	Normal	0.240	0.188	0.235	0.162
equal MSE	Normal	0.022	0.006	0.086	0.043
bias, uncond.	bootstrap	0.119	0.114	0.120	0.114
bias, condit.	bootstrap	0.118	0.115	0.132	0.110
M-Z efficiency, uncond.	bootstrap	0.085	0.061	0.116	0.075
M-Z efficiency, condit.	bootstrap	0.114	0.089	0.117	0.090
F-W efficiency, condit.	bootstrap	0.085	0.064	0.110	0.074
equal MSE	bootstrap	0.074	0.046	0.115	0.079

Notes:

1. The data generating process is a bivariate VAR(1), with coefficients given in Table 1 and error variance matrix given in section 4.1.
2. For each artificial data set, forecasts of $y_{1,t+1}$ and $y_{1,t+2}$ are formed recursively using OLS estimates of a bivariate VAR(1). We consider both unconditional forecasts and conditional forecasts obtained under the minimum-MSE approach. The conditional forecasts of $y_{1,t+1}$ and $y_{1,t+2}$ are based on a condition of $\hat{y}_{2,t,\tau}^c = y_{2,t+\tau}$, $\tau = 1, 2$. These forecasts are then used to form bias, efficiency, and accuracy tests, detailed in sections 3 and 4.2.
3. R and P refer to the number of in-sample observations and forecasts, respectively.
4. In each Monte Carlo replication, the simulated test statistics are compared against Gaussian critical values and critical values obtained with the bootstrap of the VARs described in section 3.5.
5. The number of Monte Carlo simulations is 2000; the number of bootstrap draws is 499.

Table 3: Monte Carlo Results on Size, Minimum-MSE Conditioning, DGP 2
(nominal size = 10%)

<i>test</i>	<i>source of critical values</i>	1-step horizon, variable 1				1-step horizon, variable 2			
		<i>R=50</i> <i>P=100</i>	<i>R=50</i> <i>P=150</i>	<i>R=100</i> <i>P=50</i>	<i>R=100</i> <i>P=100</i>	<i>R=50</i> <i>P=100</i>	<i>R=50</i> <i>P=150</i>	<i>R=100</i> <i>P=50</i>	<i>R=100</i> <i>P=100</i>
bias, uncond.	Normal	0.145	0.129	0.130	0.132	0.139	0.133	0.136	0.134
bias, condit.	Normal	0.132	0.132	0.122	0.124	0.150	0.143	0.144	0.134
M-Z efficiency, uncond.	Normal	0.414	0.427	0.215	0.248	0.251	0.256	0.199	0.189
M-Z efficiency, condit.	Normal	0.482	0.499	0.232	0.284	0.320	0.344	0.228	0.230
F-W efficiency, condit.	Normal	0.403	0.407	0.234	0.240	0.234	0.251	0.196	0.193
equal MSE	Normal	0.035	0.018	0.066	0.038	0.143	0.155	0.111	0.103
bias, uncond.	bootstrap	0.120	0.104	0.127	0.101	0.109	0.091	0.123	0.103
bias, condit.	bootstrap	0.119	0.118	0.124	0.107	0.111	0.096	0.133	0.101
M-Z efficiency, uncond.	bootstrap	0.096	0.086	0.107	0.099	0.097	0.102	0.104	0.108
M-Z efficiency, condit.	bootstrap	0.085	0.066	0.100	0.089	0.094	0.105	0.100	0.103
F-W efficiency, condit.	bootstrap	0.087	0.080	0.099	0.096	0.097	0.102	0.093	0.110
equal MSE	bootstrap	0.079	0.048	0.110	0.090	0.054	0.043	0.086	0.069
		2-step horizon, variable 1				2-step horizon, variable 2			
bias, uncond.	Normal	0.141	0.131	0.143	0.125	0.142	0.133	0.150	0.135
bias, condit.	Normal	0.124	0.127	0.141	0.131	0.142	0.133	0.154	0.128
M-Z efficiency, uncond.	Normal	0.312	0.292	0.211	0.206	0.315	0.310	0.294	0.242
M-Z efficiency, condit.	Normal	0.399	0.393	0.245	0.251	0.394	0.388	0.310	0.290
F-W efficiency, condit.	Normal	0.305	0.288	0.251	0.214	0.290	0.289	0.296	0.241
equal MSE	Normal	0.060	0.037	0.093	0.059	0.073	0.066	0.093	0.079
bias, uncond.	bootstrap	0.116	0.107	0.126	0.104	0.107	0.091	0.123	0.105
bias, condit.	bootstrap	0.115	0.116	0.122	0.113	0.108	0.097	0.126	0.098
M-Z efficiency, uncond.	bootstrap	0.090	0.089	0.104	0.097	0.088	0.098	0.102	0.100
M-Z efficiency, condit.	bootstrap	0.081	0.077	0.102	0.091	0.091	0.092	0.101	0.102
F-W efficiency, condit.	bootstrap	0.082	0.084	0.103	0.090	0.087	0.097	0.102	0.110
equal MSE	bootstrap	0.070	0.043	0.099	0.084	0.058	0.048	0.100	0.075
		4-step horizon, variable 1				4-step horizon, variable 2			
bias, uncond.	Normal	0.138	0.129	0.169	0.142	0.141	0.126	0.171	0.136
bias, condit.	Normal	0.132	0.131	0.168	0.144	0.147	0.118	0.177	0.136
M-Z efficiency, uncond.	Normal	0.313	0.274	0.285	0.217	0.561	0.554	0.462	0.418
M-Z efficiency, condit.	Normal	0.358	0.345	0.285	0.254	0.499	0.493	0.392	0.361
F-W efficiency, condit.	Normal	0.311	0.264	0.300	0.225	0.530	0.524	0.464	0.415
equal MSE	Normal	0.097	0.088	0.154	0.105	0.056	0.042	0.116	0.062
bias, uncond.	bootstrap	0.104	0.110	0.120	0.114	0.106	0.093	0.112	0.100
bias, condit.	bootstrap	0.103	0.107	0.120	0.111	0.110	0.090	0.117	0.102
M-Z efficiency, uncond.	bootstrap	0.066	0.059	0.099	0.064	0.061	0.061	0.092	0.069
M-Z efficiency, condit.	bootstrap	0.077	0.080	0.105	0.086	0.072	0.081	0.094	0.082
F-W efficiency, condit.	bootstrap	0.057	0.061	0.100	0.062	0.065	0.062	0.081	0.084
equal MSE	bootstrap	0.048	0.032	0.095	0.073	0.044	0.036	0.090	0.068

Notes:

1. The data generating process is a trivariate VAR(2), with coefficients given in Table 1 and error variance matrix given in section 4.1.
2. For each artificial data set, forecasts of $y_{1,t+\tau}$ and $y_{2,t+\tau}$ are formed recursively using OLS estimates of a trivariate VAR(2) and an iterative approach to computing multi-step forecasts. The conditional forecasts of $y_{1,t+\tau}$ and $y_{2,t+\tau}$ (for $\tau = 1, 2,$ and 4) are based on a condition of $y_{3,t,\tau}^c = y_{3,t+\tau}$, $\tau = 1, \dots, 4$.
3. See the notes to Table 2.

**Table 4: Monte Carlo Results on Size, Minimum-MSE Conditioning,
DGP 1, rolling estimation**
(nominal size = 10%)

1-step horizon					
<i>test</i>	<i>source of critical values</i>	<i>R=50 P=100</i>	<i>R=50 P=150</i>	<i>R=100 P=50</i>	<i>R=100 P=100</i>
bias, uncond.	Normal	0.018	0.009	0.099	0.057
bias, condit.	Normal	0.022	0.013	0.100	0.065
M-Z efficiency, uncond.	Normal	0.192	0.233	0.142	0.087
M-Z efficiency, condit.	Normal	0.212	0.269	0.142	0.088
F-W efficiency, condit.	Normal	0.160	0.148	0.122	0.073
equal MSE	Normal	0.026	0.013	0.059	0.037
bias, uncond.	bootstrap	0.116	0.097	0.123	0.114
bias, condit.	bootstrap	0.117	0.095	0.138	0.110
M-Z efficiency, uncond.	bootstrap	0.132	0.098	0.141	0.112
M-Z efficiency, condit.	bootstrap	0.122	0.076	0.143	0.106
F-W efficiency, condit.	bootstrap	0.128	0.076	0.132	0.103
equal MSE	bootstrap	0.104	0.091	0.134	0.103
2-step horizon					
bias, uncond.	Normal	0.013	0.004	0.112	0.058
bias, condit.	Normal	0.018	0.009	0.115	0.064
M-Z efficiency, uncond.	Normal	0.567	0.711	0.276	0.236
M-Z efficiency, condit.	Normal	0.324	0.417	0.200	0.136
F-W efficiency, condit.	Normal	0.435	0.508	0.245	0.195
equal MSE	Normal	0.040	0.017	0.090	0.050
bias, uncond.	bootstrap	0.118	0.101	0.126	0.117
bias, condit.	bootstrap	0.116	0.088	0.132	0.112
M-Z efficiency, uncond.	bootstrap	0.079	0.046	0.115	0.066
M-Z efficiency, condit.	bootstrap	0.099	0.059	0.126	0.095
F-W efficiency, condit.	bootstrap	0.079	0.038	0.099	0.077
equal MSE	bootstrap	0.088	0.061	0.113	0.088

Notes:

1. In these experiments, forecasts of $y_{1,t+1}$ and $y_{1,t+2}$ are formed recursively using OLS estimates of a bivariate VAR(1).
2. See the notes to Table 2.

Table 5: Monte Carlo Results on Size, Minimum-MSE Conditioning, DGP 1G
(nominal size = 10%)

1-step horizon					
<i>test</i>	<i>source of critical values</i>	<i>R=50</i> <i>P=100</i>	<i>R=50</i> <i>P=150</i>	<i>R=100</i> <i>P=50</i>	<i>R=100</i> <i>P=100</i>
bias, uncond.	Normal	0.118	0.120	0.114	0.114
bias, condit.	Normal	0.117	0.112	0.118	0.113
M-Z efficiency, uncond.	Normal	0.147	0.141	0.140	0.117
M-Z efficiency, condit.	Normal	0.192	0.173	0.155	0.147
F-W efficiency, condit.	Normal	0.134	0.130	0.134	0.122
equal MSE	Normal	0.024	0.013	0.082	0.045
bias, uncond.	bootstrap	0.109	0.124	0.129	0.108
bias, condit.	bootstrap	0.120	0.114	0.127	0.101
M-Z efficiency, uncond.	bootstrap	0.110	0.114	0.135	0.103
M-Z efficiency, condit.	bootstrap	0.117	0.102	0.120	0.097
F-W efficiency, condit.	bootstrap	0.114	0.102	0.128	0.108
equal MSE	bootstrap	0.114	0.099	0.122	0.114
2-step horizon					
bias, uncond.	Normal	0.118	0.111	0.126	0.116
bias, condit.	Normal	0.113	0.105	0.130	0.110
M-Z efficiency, uncond.	Normal	0.267	0.246	0.237	0.205
M-Z efficiency, condit.	Normal	0.227	0.199	0.218	0.180
F-W efficiency, condit.	Normal	0.224	0.223	0.217	0.183
equal MSE	Normal	0.028	0.021	0.095	0.061
bias, uncond.	bootstrap	0.110	0.123	0.130	0.108
bias, condit.	bootstrap	0.115	0.115	0.124	0.107
M-Z efficiency, uncond.	bootstrap	0.064	0.062	0.107	0.073
M-Z efficiency, condit.	bootstrap	0.107	0.092	0.123	0.095
F-W efficiency, condit.	bootstrap	0.074	0.068	0.106	0.076
equal MSE	bootstrap	0.079	0.057	0.106	0.083

Notes:

1. The data generating process is a bivariate VAR(1) with GARCH, with coefficients given in Table 1 and and section 4.1.
2. See the notes to Table 2.

Table 6: Monte Carlo Results on Power, Minimum-MSE Conditioning, DGP 3
(nominal size = 10%)

1-step horizon					
<i>test</i>	<i>source of critical values</i>	<i>R=50 P=100</i>	<i>R=50 P=150</i>	<i>R=100 P=50</i>	<i>R=100 P=100</i>
bias, uncond.	bootstrap	0.628	0.642	0.740	0.850
bias, condit.	bootstrap	0.444	0.481	0.587	0.694
M-Z efficiency, uncond.	bootstrap	0.244	0.315	0.113	0.158
M-Z efficiency, condit.	bootstrap	0.134	0.126	0.120	0.100
F-W efficiency, condit.	bootstrap	0.136	0.159	0.089	0.085
equal MSE	bootstrap	0.115	0.104	0.132	0.119
2-step horizon					
bias, uncond.	bootstrap	0.632	0.657	0.738	0.850
bias, condit.	bootstrap	0.510	0.535	0.643	0.748
M-Z efficiency, uncond.	bootstrap	0.230	0.314	0.108	0.173
M-Z efficiency, condit.	bootstrap	0.104	0.113	0.099	0.079
F-W efficiency, condit.	bootstrap	0.115	0.127	0.079	0.071
equal MSE	bootstrap	0.104	0.064	0.171	0.149

Notes:

1. The data generating process is a bivariate VAR(1), with coefficient breaks, using the coefficient values given in Table 1 and the error variance matrix given in section 4.1. In each experiment, the coefficient break occurs in period $R+1$.
2. See the notes to Table 2.

Table 7: Monte Carlo Results on Power, Minimum-MSE Conditioning, DGP 4
(nominal size = 10%)

1-step horizon					
<i>test</i>	<i>source of critical values</i>	<i>R=50 P=100</i>	<i>R=50 P=150</i>	<i>R=100 P=50</i>	<i>R=100 P=100</i>
bias, uncond.	bootstrap	0.141	0.120	0.122	0.106
bias, condit.	bootstrap	0.096	0.103	0.097	0.069
M-Z efficiency, uncond.	bootstrap	0.129	0.104	0.128	0.114
M-Z efficiency, condit.	bootstrap	0.199	0.119	0.609	0.466
F-W efficiency, condit.	bootstrap	0.147	0.103	0.169	0.178
equal MSE	bootstrap	0.490	0.398	0.945	0.876
2-step horizon					
bias, uncond.	bootstrap	0.153	0.140	0.128	0.115
bias, condit.	bootstrap	0.091	0.103	0.099	0.072
M-Z efficiency, uncond.	bootstrap	0.082	0.067	0.127	0.086
M-Z efficiency, condit.	bootstrap	0.335	0.198	0.757	0.723
F-W efficiency, condit.	bootstrap	0.112	0.081	0.147	0.130
equal MSE	bootstrap	0.459	0.418	0.854	0.822

Notes:

1. The data generating process is a bivariate VAR(1), with a break in the error correlation described in section 4.1. In each experiment, the coefficient break occurs in period $R+1$.
2. See the notes to Table 2.

Table 8: Monte Carlo Results on Size and Power, Policy Shock Conditioning
(nominal size = 10%)

DGP 1, 2-step horizon					
<i>test</i>	<i>source of critical values</i>	<i>R=50</i>	<i>R=50</i>	<i>R=100</i>	<i>R=100</i>
		<i>P=100</i>	<i>P=150</i>	<i>P=50</i>	<i>P=100</i>
bias	Normal	0.118	0.126	0.128	0.123
M-Z efficiency	Normal	0.176	0.147	0.207	0.133
F-W efficiency	Normal	0.256	0.229	0.251	0.173
bias	bootstrap	0.118	0.116	0.126	0.112
M-Z efficiency	bootstrap	0.074	0.056	0.119	0.075
F-W efficiency	bootstrap	0.083	0.057	0.117	0.066
DGP 3, 2-step horizon					
bias	bootstrap	0.627	0.652	0.732	0.842
M-Z efficiency	bootstrap	0.179	0.243	0.098	0.141
F-W efficiency	bootstrap	0.189	0.238	0.093	0.130
DGP 4, 2-step horizon					
bias	bootstrap	0.142	0.123	0.131	0.106
M-Z efficiency	bootstrap	0.085	0.061	0.126	0.115
F-W efficiency	bootstrap	0.088	0.073	0.117	0.088

Notes:

1. In each experiment, the data generating process is a bivariate VAR(1), with coefficients given in Table 1 and error variance matrix given in section 4.1.
2. For each artificial data set, forecasts of $y_{1,t+1}$ and $y_{1,t+2}$ are formed recursively using OLS estimates of a bivariate VAR(1). We consider both unconditional forecasts and conditional forecasts obtained under the policy shock approach. The conditional forecasts of $y_{1,t+1}$ and $y_{1,t+2}$ are based on a condition of $\hat{y}_{2,t,\tau}^c = y_{2,t+\tau}$, $\tau = 1, 2$. These forecasts are then used to form bias, efficiency, and accuracy tests, detailed in sections 3 and 4.2. Since the 1-step ahead forecasts are equivalent to the unconditional forecast, we only report results for the 2-step ahead horizon.
3. See the notes to Table 2.

Table 9. Tests of unconditional and conditional (min.-MSE) forecasts from 22-variable BVAR, 1991-2007

(significant tests bolded; 10% bootstrap critical values in parentheses)

GDP growth	$\tau=1$	$\tau=2$	$\tau=4$	$\tau=8$
bias, uncond.	-1.279 (-1.450, 0.585)	-1.306 (-1.654, 0.537)	-0.910 (-1.612, 0.761)	-0.298 (-2.242, 1.439)
bias, condit.	-0.827 (-1.261, 0.534)	-0.731 (-1.462, 0.737)	-0.641 (-1.609, 0.957)	-0.439 (-2.381, 1.192)
M-Z efficiency, uncond.	-2.774 (-1.463, 1.727)	-3.447 (-2.416, 1.793)	-4.241 (-3.621, 1.216)	-5.144 (-5.107, 1.259)
M-Z efficiency, condit.	-3.714 (-2.059, 1.510)	-3.753 (-2.785, 1.576)	-2.878 (-2.738, 2.354)	-5.736 (-5.690, 2.051)
F-W efficiency, condit.	-3.049 (-1.674, 1.767)	-3.961 (-2.804, 1.862)	-4.933 (-3.615, 1.224)	-5.460 (-4.514, 1.583)
Unemployment rate	$\tau=1$	$\tau=2$	$\tau=4$	$\tau=8$
bias, uncond.	0.545 (-1.071, 2.288)	0.634 (-0.935, 2.130)	0.618 (-0.960, 1.980)	0.435 (-0.989, 1.949)
bias, condit.	0.069 (-0.837, 1.780)	0.158 (-0.797, 1.710)	0.307 (-0.849, 1.698)	0.323 (-0.996, 1.767)
M-Z efficiency, uncond.	0.987 (-2.282, 1.024)	0.635 (-2.502, 1.148)	0.156 (-3.316, 1.055)	-1.642 (-6.429, 0.938)
M-Z efficiency, condit.	1.638 (-2.586, 1.064)	1.197 (-2.643, 1.212)	0.758 (-3.718, 1.357)	-0.890 (-6.993, 1.262)
F-W efficiency, condit.	1.561 (-2.693, 1.030)	1.201 (-2.970, 1.287)	0.829 (-4.272, 1.178)	-1.074 (-7.283, 0.840)
Core PCE inflation	$\tau=1$	$\tau=2$	$\tau=4$	$\tau=8$
bias, uncond.	-2.166 (-2.820, 1.233)	-2.139 (-2.919, 1.192)	-2.326 (-3.198, 1.418)	-2.630 (-3.458, 1.911)
bias, condit.	-1.962 (-2.835, 1.417)	-2.065 (-2.847, 1.371)	-1.885 (-2.864, 1.523)	-1.460 (-3.164, 2.146)
M-Z efficiency, uncond.	-4.569 (-3.688, 1.003)	-4.140 (-4.048, 0.616)	-5.423 (-5.006, 0.589)	-8.316 (-7.851, 1.007)
M-Z efficiency, condit.	-4.076 (-3.805, 1.305)	-4.044 (-4.484, 0.707)	-6.294 (-5.362, 0.526)	-18.110 (-8.101, 0.661)
F-W efficiency, condit.	-3.945 (-3.673, 1.093)	-4.156 (-4.374, 0.731)	-5.612 (-5.367, 0.715)	-7.848 (-7.677, 1.057)

Notes:

1. As described in section 5.1, forecasts of real GDP growth, the unemployment rate, and core PCE inflation (all defined at annualized rates) are obtained from recursive estimates of a 22-variable BVAR. The forecasts included are unconditional and minimum-MSE conditional. At each forecast origin t , the conditions imposed are that, over an eight quarter forecast horizon from $t + 1$ through $t + 8$, the federal funds rate take its actual values over the period.
2. The bias, efficiency, and MSE accuracy tests detailed in section 4.2 are compared against standard normal critical values and critical values obtained with the bootstrap described in section 3.5. The number of bootstrap draws is 999.

Table 10. Accuracy of unconditional and conditional (min.-MSE) forecasts from 22-variable BVAR, 1991-2007

(significant tests bolded; 10% bootstrap critical values in parentheses)

GDP growth	$\tau=1$	$\tau=2$	$\tau=4$	$\tau=8$
unconditional MSE (RMSE)	3.407 (1.846)	3.977 (1.994)	4.485 (2.118)	5.155 (2.270)
conditional MSE (RMSE)	3.546 (1.883)	3.953 (1.988)	3.650 (1.910)	5.869 (2.423)
$MSE_U - MSE_C$	-0.139	0.024	0.835	-0.714
population estimate of $MSE_U - MSE_C$	0.019	0.088	0.322	0.808
MSE <i>t</i> -test	-0.672	-0.206	1.085	-1.749
10% bootstrap crit. vals.	(-0.846, 1.941)	(-0.448, 2.603)	(0.288, 4.102)	(-3.095, 2.728)
Unemployment rate	$\tau=1$	$\tau=2$	$\tau=4$	$\tau=8$
unconditional MSE (RMSE)	0.020 (0.142)	0.052 (0.228)	0.197 (0.444)	0.704 (0.839)
conditional MSE (RMSE)	0.018 (0.133)	0.041 (0.203)	0.137 (0.370)	0.455 (0.675)
$MSE_U - MSE_C$	0.002	0.011	0.060	0.249
population estimate of $MSE_U - MSE_C$	0.007	0.028	0.094	0.204
MSE <i>t</i> -test	-2.046	-1.968	-0.796	0.356
10% bootstrap crit. vals.	(-2.898, 1.718)	(-2.046, 1.681)	(-1.315, 1.982)	(-1.127, 2.169)
Core PCE inflation	$\tau=1$	$\tau=2$	$\tau=4$	$\tau=8$
unconditional MSE (RMSE)	0.316 (0.562)	0.435 (0.660)	0.500 (0.707)	0.930 (0.964)
conditional MSE (RMSE)	0.291 (0.539)	0.373 (0.611)	0.478 (0.691)	1.112 (1.054)
$MSE_U - MSE_C$	0.025	0.062	0.022	-0.182
population estimate of $MSE_U - MSE_C$	0.001	0.007	0.050	0.403
MSE <i>t</i> -test	1.615	1.310	-0.390	-4.287
10% bootstrap crit. vals.	(-1.316, 2.178)	(-1.156, 2.183)	(-1.363, 2.037)	(-3.764, 1.816)

Notes:

1. The table reports forecast MSEs (and RMSEs) for the 1991-2007 sample. It also includes estimates of the population difference in MSEs (unconditional less conditional) based on the full sample VAR estimates, computed as described in section 3.4.
2. See the notes to Table 9.